

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 October 2002 (10.10.2002)

PCT

(10) International Publication Number
WO 02/079169 A1

(51) International Patent Classification⁷: C07D 239/42,
409/04, 401/04, 403/12, A61K 31/505, 31/506, A61P 9/14

(21) International Application Number: PCT/US02/09494

(22) International Filing Date: 26 March 2002 (26.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/279,956 29 March 2001 (29.03.2001) US
Not furnished 22 March 2002 (22.03.2002) US

(71) Applicant (for all designated States except US): BRIS-
TOL-MYERS SQUIBB COMPANY [US/US]; P.O. BOX
4000, Rt. 206 & Province Line Road, Princeton, NJ 08543-
4000 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LOMBARDO,
Louis, J. [US/US]; 49 West Street, Belle Mead, NJ 08502
(US). WU, Laurence, I [US/US]; 43 Nathan Court,
Newtown, PA 18940 (US).

(74) Agents: GIBBONS, Maureen et al.; Bristol-Myers
Squibb Company, P.O. Box 4000, Rt. 206 & Province Line
Road, Princeton, NJ 08543-4000 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).


Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments
- entirely in electronic form (except for this front page) and
available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 02/079169 A1

(54) Title: CYANO-SUBSTITUTED DIHYDROPYRIMIDINE COMPOUNDS AND THEIR USE TO TREAT DISEASES



(I)

(57) Abstract: The present invention provides compounds of formula (I), formula (I) and pharmaceutically acceptable salts thereof. The formula I compounds induce mitotic arrest thereby making them useful as anti-cancer agents. The formula I compounds are also useful for the treatment of other diseases which can be treated by inducing mitotic arrest.

BEST AVAILABLE COPY

CYANO-SUBSTITUTED DIHYDROPYRIMIDINE COMPOUNDS AND THEIR USE TO TREAT DISEASES

RELATED APPLICATIONS

5 This application claims the benefit under 35 U.S.C. Section 119(e) of U.S.
Provisional Patent Application No. 60/279,956 filed March 29, 2001.

FIELD OF INVENTION

10 This invention relates to novel compounds that interrupt mitosis thereby
making the compounds useful for the treatment of proliferative diseases, such as
cancer.

BACKGROUND

15 Cell proliferation and programmed cell death play important roles in the
growth and development of an organism. In proliferative diseases such as cancer, the
processes of cell proliferation and/or programmed cell death are often perturbed. For
example, a cancer cell may have unregulated cell division through either the
overexpression of a positive regulator of the cell cycle or the loss of a negative
regulator of the cell cycle, perhaps by mutation. Alternatively, a cancer cell may have
20 lost the ability to undergo programmed cell death through the overexpression of a
negative regulator of apoptosis. Hence, there is a need to develop new
chemotherapeutic drugs that will restore the processes of checkpoint control and
programmed cell death to cancerous cells.

25 One approach to the treatment of human cancers is to target a protein that is
essential for cell cycle progression. In order for the cell cycle to proceed from one
phase to the next, certain prerequisite events must be completed. There are
checkpoints within the cell cycle that enforce the proper order of events and phases.
One such checkpoint is the spindle checkpoint that occurs during the metaphase stage
of mitosis. Small molecules that target proteins with essential functions in mitosis
30 may initiate the spindle checkpoint to arrest cells in mitosis. Of the small molecules
that arrest cells in mitosis, those which display anti-tumor activity in the clinic also
induce apoptosis, the morphological changes associated with programmed cell death.

An effective chemotherapeutic for the treatment of cancer may be one that induces checkpoint control and subsequent programmed cell death.

Most compounds known to cause mitotic arrest and apoptosis act as tubulin binding agents. These compounds alter the dynamic instability of microtubules and indirectly alter the function/structure of the mitotic spindle thereby causing mitotic arrest. Because most of these compounds target the tubulin protein, a component of all microtubules, they may also affect normal cellular processes in which microtubules have a role. Hence, a need exists for small molecules that specifically target proteins associated with proliferating cells, such as Eg5.


Eg5 is one of several kinesin-like motor proteins that are localized to the mitotic spindle and known to be required for formation and/or function of the bipolar mitotic spindle. Recently, there was a report of a small molecule that disturbs bipolarity of the mitotic spindle (Mayer, T.U. et. al. 1999. Science 286(5441) 971-4). More specifically, the small molecule induced the formation of an aberrant mitotic spindle wherein a monoastral array of microtubules emanated from a central pair of centrosomes, with chromosomes attached to the distal ends of the microtubules. The small molecule was dubbed "monastrol" after the monoastral array. This monoastral array phenotype had been previously observed in mitotic cells that were immunodepleted of the Eg5 motor protein.

The distinctive monoastral array phenotype facilitated identification of monastrol as a potential inhibitor of Eg5. Indeed, monastrol was further shown to inhibit the Eg5 motor-driven motility of microtubules in an *in vitro* assay. Furthermore, monastrol had no apparent effect upon the related kinesin motor or upon the motor(s) responsible for golgi apparatus movement within the cell. Cells that display the monoastral array phenotype, either through immunodepletion of Eg5 or monastrol inhibition of Eg5, arrest in M-phase of the cell cycle. Unfortunately, however, the mitotic arrest induced by either of these mechanisms is transient. (Kapoor, 2000. *J. Cell. Biol.* 150(5) 975-80). Both the monoastral array phenotype and the monastrol induced cell cycle arrest in mitosis are reversible. Cells recover to form a normal bipolar mitotic spindle, to complete mitosis, and to proceed through the cell cycle and normal cell proliferation. This suggests that a small molecule inhibitor of Eg5 that induced a transient mitotic arrest may not be effective for the treatment of

cancer cell proliferation. Nonetheless, the discovery that monastrol causes mitotic arrest is intriguing and hence there is a need to further study and identify compounds that can be used to modulate the Eg5 motor protein in a manner that would be effective in the treatment of human cancers. There is also a need to explore the use of these compounds in combination with other antineoplastic agents.

SUMMARY

The compounds of the invention cause the interruption of mitosis, and as such, can be used to treat proliferative diseases. For example, the compounds of the instant invention can be used as antiproliferatives and anticancer agents. More specifically, the invention comprises a compound of formula I



its enantiomers, diastereomers, pharmaceutically acceptable salts, prodrugs and solvates thereof, wherein

R_1 is selected from the group consisting of hydrogen, alkyl and cycloalkyl;

R_2 and R_3 are each independently selected from the group consisting of H, alkyl, aryl, heteroaryl, arylalkyl, cycloalkyl, heterocycloalkyl, cycloalkylalkyl, heterocycloalkylalkyl and heteroarylalkyl; or

R_2 and R_3 may also be taken together to form a carbocyclic or heterocyclic ring;

R_4 is selected from the group consisting of alkyl, arylalkyl, cycloalkylalkyl, heteroarylalkyl, heterocycloalkyl, aminoalkyl, heterocycloalkylalkyl, CN, $C(O)R_5$, CO_2R_5 , $C(O)SR_5$ and $CONR_5R_6$;


R_5 and R_6 are each independently selected from the group consisting of H, alkyl, cycloalkyl, hydroxyalkyl, alkenyl, alkoxy, thioalkoxy, alkoxyalkyl, haloalkyl, aryl, heteroaryl, arylalkyl, cycloalkylalkyl, heteroarylalkyl and heterocycloalkylalkyl or $N-R_5R_6$ together form a heterocycloalkyl;

Z is selected from the group consisting of O, S and NR_8 ;

R_8 is selected from the group consisting of H, CN, sulfonamido, OR_7 , alkyl, cycloalkyl, aryl, arylalkyl, heterocyclyl, heteroaryl and heteroarylalkyl; and

R_7 is selected from the group consisting of H, alkyl, arylalkyl, cycloalkylalkyl, heterocycloalkylalkyl and heteroarylalkyl.

5 According to one embodiment, compounds of the present invention include those having formula IA:



10

its enantiomers, diastereomers, pharmaceutically acceptable salts, prodrugs and solvates thereof, wherein Y_1 , Y_2 , Y_3 , Y_4 and Y_5 are independently selected from the group consisting of H, halogen, lower alkyl, O-alkyl, O-aryl, NH-alkyl, NH-aryl, N-alkylalkyl, N-alkylaryl, N-arylaryl with the proviso that at least one of Y_1 , Y_2 , Y_3 , Y_4 and Y_5 is aryl or heteroaryl, and the remaining groups are as defined above.

The present invention also provides methods for treating a proliferative disease, such as cancer, via modulation of the Eg5 motor protein comprising administering to a mammalian species in need of such treatment an effective amount of at least one compound of formula I or IA, as defined above.

20

DESCRIPTION

The present invention provides for compounds of formula I and IA, as defined above, pharmaceutical compositions employing such compounds, and methods of using such compounds.

25

Listed below are definitions of various terms used to describe the compounds of the instant invention. These definitions apply to the terms as they are used

throughout the specification (unless they are otherwise limited in specific instances) either individually or as part of a larger group.

The term "alkyl" herein alone or as part of another group refers to a monovalent alkane (hydrocarbon) derived radical containing from 1 to 12 carbon atoms unless otherwise defined. An alkyl group is an optionally substituted straight, branched or cyclic saturated hydrocarbon group. When substituted, alkyl groups may be substituted with up to four substituent groups, R as defined, at any available point of attachment. When the alkyl group is said to be substituted with an alkyl group, this is used interchangeably with "branched alkyl group". Exemplary unsubstituted such groups include methyl, ethyl, propyl, isopropyl, n-butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl, 2,2,4-trimethylpentyl, nonyl, decyl, undecyl, dodecyl, and the like. Exemplary substituents may include but are not limited to one or more of the following groups: aryl, halo (such as F, Cl, Br, I), haloalkyl (such as CCl₃ or CF₃), alkoxy, alkylthio, hydroxy, carboxy (-COOH), alkoxycarbonyl (-C(O)R), alkylcarbonyloxy (-OCOR), amino (-NH₂), carbamoyl (-NHCOOR- or -OCONHR-), urea (-NHCONHR-) or thiol (-SH). Alkyl groups as defined may also comprise one or more carbon to carbon double bonds or one or more carbon to carbon triple bonds.

The term "alkenyl" herein alone or as part of another group refers to a hydrocarbon radical straight, branched or cyclic containing from 2 to 12 carbon atoms and at least one carbon to carbon double bond.

The term "alkynyl" herein alone or as part of another group refers to a hydrocarbon radical straight, branched or cyclic containing from 2 to 12 carbon atoms and at least one carbon to carbon triple bond.

The numbers in the subscript after the symbol "C" define the number of carbon atoms a particular group can contain. For example "C₁₋₆ alkyl" means a straight or branched saturated carbon chain having from one to six carbon atoms; examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, t-butyl, n-pentyl, sec-pentyl, isopentyl, and n-hexyl. Depending on the context, "C₁₋₆ alkyl" can also refer to C₁₋₆ alkylene which bridges two groups; examples include propane-1,3-diyl, butane-1,4-diyl, 2-methyl-butane-1,4-diyl, etc. "C₂₋₆ alkenyl" means a straight or branched carbon chain having at least one carbon-carbon double

bond, and having from two to six carbon atoms; examples include ethenyl, propenyl, isopropenyl, butenyl, isobutenyl, pentenyl, and hexenyl. Depending on the context, "C₂₋₆ alkenyl" can also refer to C₂₋₆ alkenediyl which bridges two groups; examples include ethylene-1,2-diyl (vinylene), 2-methyl-2-butene-1,4-diyl, 2-hexene-1,6-diyl, etc. "C₂₋₆ alkynyl" means a straight or branched carbon chain having at least one carbon-carbon triple bond, and from two to six carbon atoms; examples include ethynyl, propynyl, butynyl, and hexynyl.

The term "cycloalkyl" herein alone or as part of another group is a species of alkyl containing from 3 to 15 carbon atoms, without alternating or resonating double bonds between carbon atoms. It may contain from 1 to 4 rings. Exemplary unsubstituted such groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, adamantyl, etc. Exemplary substituents include one or more of the following groups: halogen, alkyl, alkoxy, alkyl hydroxy, amino, nitro, cyano, thiol and/or alkylthio.

The terms "alkoxy" or "alkylthio" herein alone or as part of another group denote an alkyl group as described above bonded through an oxygen linkage (-O-) or a sulfur linkage (-S-), respectively.

The term "alkyloxycarbonyl" herein alone or as part of another group denotes an alkoxy group bonded through a carbonyl group. An alkoxy carbonyl radical is represented by the formula: -C(O)OR, where the R group is a straight or branched C₁₋₆ alkyl group.

The term "alkylcarbonyl" herein alone or as part of another group refers to an alkyl group bonded through a carbonyl group.

The term "alkylcarbonyloxy" herein alone or as part of another group denotes an alkylcarbonyl group which is bonded through an oxygen linkage.

The term "arylalkyl" herein alone or as part of another group denotes an aromatic ring bonded to an alkyl group as described above.

The term "aryl" herein alone or as part of another group refers to monocyclic or bicyclic aromatic rings, e.g. phenyl, substituted phenyl and the like, as well as groups which are fused, e.g., naphthyl, phenanthrenyl and the like. An aryl group thus contains at least one ring having at least 6 atoms, with up to five such rings being present, containing up to 22 atoms therein, with alternating (resonating) double bonds between adjacent carbon atoms or suitable heteroatoms. Aryl groups may optionally

be substituted with one or more groups including, but not limited to halogen, alkyl, alkoxy, hydroxy, carboxy, carbamoyl, alkylloxycarbonyl, nitro, alkenyloxy, trifluoromethyl, amino, cycloalkyl, aryl, heteroaryl, cyano, alkyl S(O)_m (m=0, 1, 2), or thiol.

5 The term "amino" herein alone or as part of another group refers to -NH₂. An "amino" may optionally be substituted with one or two substituents, which may be the same or different, such as alkyl, aryl, arylalkyl, alkenyl, alkynyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, thioalkyl, carbonyl or carboxyl. These
10 substituents may be further substituted with a carboxylic acid, any of the alkyl or aryl substituents set out herein. In some embodiments, the amino groups are substituted with carboxyl or carbonyl to form N-acyl or N-carbamoyl derivatives

 The term "carbocyclic ring" herein alone or as part of another group refers to stable, saturated or partially unsaturated monocyclic ring hydrocarbyls of 3 to 7
15 carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. The carbocyclic ring may be optionally substituted meaning that the carbocyclic ring may be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower alkyl), alkoxy (preferably lower alkoxy), nitro, monoalkylamino (preferably a lower alkylamino),
20 dialkylamino (preferably a di[lower]alkylamino), cyano, halo, haloalkyl (preferably trifluoromethyl), alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl
25 (preferably phenyl), said aryl being optionally substituted by halo, lower alkyl and lower alkoxy groups.

 The term "cycloalkyl" herein alone or as part of another group refers to fully saturated and partially unsaturated hydrocarbon rings of 3 to 9, preferably 3 to 7 carbon atoms. Further, a cycloalkyl may be substituted. A substituted cycloalkyl
30 refers to such rings having one, two, or three substituents, preferably one, selected from the group consisting of halo, alkyl, substituted alkyl, alkenyl, alkynyl, nitro, cyano, oxo (=O), hydroxy, alkoxy, thioalkyl, -CO₂H, -C(=O)H, CO₂-alkyl, -

C(=O)alkyl, keto, =N-OH, =N-O-alkyl, aryl, heteroaryl, heterocyclo, a five or six membered ketal (*i.e.* 1,3-dioxolane or 1,3-dioxane), -NR'R'', -C(=O)NR'R'', -CO₂NR'R'', -C(=O)NR'R'', -NR'CO₂R'', -NR'C(=O)R'', -SO₂NR'R'', and -NR'SO₂R'', wherein each of R' and R'' is independently selected from hydrogen, alkyl, substituted alkyl, and cycloalkyl, or R' and R'' together form a heterocyclo or heteroaryl ring.

The term "heteroaryl" herein alone or as part of another group refers to substituted and unsubstituted aromatic 5 or 6 membered monocyclic groups, 9 or 10 membered bicyclic groups, and 11 to 14 membered tricyclic groups which have at least one heteroatom (O, S or N) in at least one of the rings. Each ring of the heteroaryl group containing a heteroatom can contain one or two oxygen or sulfur atoms and/or from one to four nitrogen atoms provided that the total number of heteroatoms in each ring is four or less and each ring has at least one carbon atom. The fused rings completing the bicyclic and tricyclic groups may contain only carbon atoms and may be saturated, partially saturated, or unsaturated. The nitrogen and sulfur atoms may optionally be oxidized and the nitrogen atoms may optionally be quaternized. Heteroaryl groups which are bicyclic or tricyclic must include at least one fully aromatic ring but the other fused ring or rings may be aromatic or non-aromatic. The heteroaryl group may be attached at any available nitrogen or carbon atom of any ring. The heteroaryl ring system may contain zero, one, two or three substituents selected from the group consisting of halo, alkyl, substituted alkyl, alkenyl, alkynyl, aryl, nitro, cyano, hydroxy, alkoxy, thioalkyl, -CO₂H, -C(=O)H, -CO₂-alkyl, -C(=O)alkyl, phenyl, benzyl, phenylethyl, phenyloxy, phenylthio, cycloalkyl, substituted cycloalkyl, heterocyclo, heteroaryl, -NR'R'', -C(=O)NR'R'', -CO₂NR'R'', -C(=O)NR'R'', -NR'CO₂R'', -NR'C(=O)R'', -SO₂NR'R'', and -NR'SO₂R'', wherein each of R' and R'' is independently selected from hydrogen, alkyl, substituted alkyl, and cycloalkyl, or R' and R'' together form a heterocyclo or heteroaryl ring.

Exemplary monocyclic heteroaryl groups include pyrrolyl, pyrazolyl, pyrazolinyl, imidazolyl, oxazolyl, diazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl, furanyl, thienyl, oxadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl and the like.

Exemplary bicyclic heteroaryl groups include indolyl, benzothiazolyl, benzodioxolyl, benzoxazolyl, benzothienyl, quinolinyl, tetrahydroisoquinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, indolizynyl, benzofuranyl, chromonyl, coumarinyl, benzopyranyl, cinnolinyl, quinoxalinyl, indazolyl, pyrrolopyridyl, furopyridinyl, dihydroisoindolyl, tetrahydroquinolinyl and the like.

Exemplary tricyclic heteroaryl groups include carbazolyl, benzidolyl, phenanthrollinyl, acridinyl, phenanthridinyl, xanthenyl and the like.

The term "heterocycloalkyl" herein alone or as part of another group refers to a cycloalkyl group (nonaromatic) in which one of the carbon atoms in the ring is replaced by a heteroatom selected from O, S or N, and in which up to three additional carbon atoms may be replaced by said heteroatoms.

The term "heterocyclic ring" herein alone or as part of another group refers to a stable, saturated, or partially unsaturated monocyclic ring system containing 5 to 7 ring members of carbon atoms and other atoms selected from nitrogen, sulfur and/or oxygen. Preferably, a heterocyclyl is a 5 or 6-membered monocyclic ring and contains one, two, or three heteroatoms selected from nitrogen, oxygen and/or sulfur. The heterocyclic ring may be optionally substituted which means that the heterocyclic ring may be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower alkyl), alkoxy (preferably lower alkoxy), nitro, monoalkylamino (preferably a lower alkylamino), dialkylamino (preferably a di[lower]alkylamino), cyano, halo, haloalkyl (preferably trifluoromethyl), alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl (preferably phenyl), said aryl being optionally substituted by halo, lower alkyl and lower alkoxy groups. Examples of such heterocyclic rings are isoxazolyl, imidazolynyl, thiazolynyl, imidazolidinyl, pyrrolyl, pyrrolinyl, pyranyl, pyrazinyl, piperidyl, morpholinyl and triazolyl. The heterocyclic ring may be attached to the parent structure through a carbon atom or through any heteroatom of the heterocyclyl that results in a stable structure.

The term "heterocyclyl" herein alone or as part of another group as used herein refers to a stable, saturated, or partially unsaturated, monocyclic, bridged monocyclic, bicyclic, and spiro ring system containing carbon atoms and other atoms selected from nitrogen, sulfur and/or oxygen. Preferably, a heterocyclyl is a 5 or 6-membered
5 monocyclic ring or an 8-11 membered bicyclic ring which consists of carbon atoms and contains one, two, or three heteroatoms selected from nitrogen, oxygen and/or sulfur. The term "optionally substituted" as it refers to "heterocyclyl" herein indicates that the heterocyclyl group may be substituted at one or more substitutable ring positions by one or more groups independently selected from alkyl (preferably lower
10 alkyl), alkoxy (preferably lower alkoxy), nitro, monoalkylamino (preferably a lower alkylamino), dialkylamino (preferably a di[lower]alkylamino), cyano, halo, haloalkyl (preferably trifluoromethyl), alkanoyl, aminocarbonyl, monoalkylaminocarbonyl, dialkylaminocarbonyl, alkyl amido (preferably lower alkyl amido), alkoxyalkyl (preferably a lower alkoxy[lower]alkyl), alkoxycarbonyl (preferably a lower
15 alkoxycarbonyl), alkylcarbonyloxy (preferably a lower alkylcarbonyloxy) and aryl (preferably phenyl); said aryl being optionally substituted by halo, lower alkyl and lower alkoxy groups. Examples of such heterocyclyl groups are isoxazolyl, imidazolyl, thiazolyl, imidazolidinyl, pyrrolyl, pyrrolinyl, pyranyl, pyrazinyl, piperidyl, morpholinyl and triazolyl. The heterocyclyl group may be attached to the
20 parent structure through a carbon atom or through any heteroatom of the heterocyclyl that results in a stable structure.

The term "heteroatom" means O, S or N, selected on an independent basis. It should be noted that any heteroatom with unsatisfied valences is assumed to have the hydrogen atom to satisfy the valences.

25 The term "halogen" or "halo" refers to chlorine, bromine, fluorine or iodine selected on an independent basis.

When a functional group is termed "protected", this means that the group is in modified form to preclude undesired side reactions at the protected site. Suitable protecting groups for the compounds of the present invention will be recognized from
30 the present application taking into account the level of skill in the art, and with reference to standard textbooks, such as Greene, T. W. et al., *Protective Groups in Organic Synthesis*, Wiley, N.Y. (1991).

As used herein, the term "patient" encompasses all mammalian species.

Suitable examples of salts of the compounds according to the invention with inorganic or organic acids are hydrochloride, hydrobromide, sulfate, methanesulfonate, maleate, fumarate, and phosphate. Salts which are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free compounds I or their pharmaceutically acceptable salts, are also included.


All stereoisomers of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure form. The definition of the compounds according to the invention embraces all possible stereoisomers and their mixtures. It very particularly embraces the racemic forms and the isolated optical isomers having the specified activity. The racemic forms can be resolved by physical methods, such as, for example, fractional crystallization, separation or crystallization of diastereomeric derivatives or separation by chiral column chromatography. The individual optical isomers can be obtained from the racemates by conventional methods, such as, for example, salt formation with an optically active acid followed by crystallization.

It should be understood that the present invention includes prodrug forms of the compounds of formula I. Various forms of prodrugs are well known in the art.

For examples of such prodrug derivatives, see:

- (a) Design of Prodrugs, edited by H. Bundgaard (Elsevier, 1985); and Methods in Enzymology, Vol. 42, pp. 309-396, edited by K. Widder et al., (Academic Press, 1985);
- (b) A Textbook of Drug Design and Development, edited by Krosgaard-Larsen and H. Bundgaard, Chapter 5, "Design and Application of Prodrugs," by H. Bundgaard, pp. 113-191 (1991);
- (c) H. Bundgaard, Advanced Drug Deliver Reviews, 8, pp. 1-38 (1992);
- (d) H. Bundgaard et al., Journal of Pharmaceutical Sciences, 77, 285 (1988); and
- (e) N. Kayeka et al., Chem. Phar. Bull., 32, 692 (1984).

In general, the instant invention comprises a compound of formula I



I,

its enantiomers, diastereomers, pharmaceutically acceptable salts, prodrugs and solvates thereof. R_1 is hydrogen, alkyl or cycloalkyl. R_2 and R_3 are each
 5 independently H, alkyl, aryl, heteroaryl, arylalkyl, cycloalkylalkyl, heterocycloalkylalkyl or heteroarylalkyl. Alternatively, R_2 and R_3 may be taken together to form either a carbocyclic or heterocyclic ring. R_4 is alkyl, arylalkyl, cycloalkylalkyl, aminoalkyl, heteroarylalkyl, heterocycloalkylalkyl, CN, COR_5 , CO_2R_5 or $CONR_5R_6$. R_5 and R_6 are each independently H, alkyl, arylalkyl,
 10 cycloalkylalkyl, heteroarylalkyl or heterocycloalkylalkyl or $N-R_5R_6$ together form a heterocycloalkyl; Z is O, S or NR_8 ; R_8 is H, CN, sulfonamido, OR_7 , alkyl, cycloalkyl, aryl, arylalkyl, heterocyclyl, heteroaryl or heteroarylalkyl. R_7 is H, alkyl, arylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, or heteroarylalkyl.

One preferred embodiment of the instant invention includes compounds of
 15 formula I, as defined above, wherein R_1 is alkyl; R_2 is selected from the group consisting of aryl and heteroaryl; R_3 is H; R_4 is selected from the group consisting of alkyl, arylalkyl, CO_2R_5 and $CONR_5R_6$; R_5 and R_6 are independently selected from the group consisting of H, alkyl, aminoalkyl, hydroxyalkyl, phenylamino and arylalkyl; Z is selected from the group consisting of O, S and NR_8 ; and R_8 is selected from the
 20 group consisting of H and CN.

In one preferred embodiment, the invention comprises compounds of formula I wherein R_2 is heteroaryl. Preferred heteroaryl groups include optionally substituted thiophenyl, oxazolyl, isoxazolyl, and furanyl. Preferred substituents include methyl, ethyl, halo, haloalkyl, such as CF_3 or aryl groups, such as phenyl or substituted
 25 phenyl. Preferred phenyl substituents include alkyl groups, such as methyl, alkoxy groups, such as methoxy, halo, (F, Cl, or Br), haloalkyl, such as, CF_3 , amino groups, such as dimethylamino, or alkoxyalkyl groups, such as ethoxymethyl groups.

In another preferred embodiment, the invention comprises compounds of formula I, as defined above, wherein R_4 is selected from the group consisting of alkyl,
 30 arylalkyl, CO_2R_5 and $CONR_5R_6$.


In yet another preferred embodiment, the instant invention comprises the compounds of formula I, as defined above, wherein R_4 is CO_2R_5 ; Z is O; and R_5 is ethyl.

In yet a further preferred embodiment, the instant invention comprises the compounds of formula I, as defined above, wherein R_4 is CONR_5R_6 ; Z is O; R_5 is H; and R_6 is methyl, ethyl, propyl, phenyl, cyclopropyl, hydroxyethyl, thiophenyl, or 2-propylene.

In still yet another preferred embodiment, the instant invention comprises the compounds of formula I, as defined above, wherein R_1 is CH_3 ; R_2 is aryl; R_4 is CO_2R_5 ; R_5 is alkyl; and Z is O.

In still yet another preferred embodiment, the instant invention comprises the compounds of formula I, as defined above, wherein R_1 is CH_3 ; R_2 is aryl; R_4 is CONR_5R_6 ; R_5 is alkyl, R_6 is H; and Z is O.

According to one embodiment, the instant invention comprises compounds of formula IA:



and include its enantiomers, diastereomers, pharmaceutically acceptable salts, prodrugs and solvates thereof, wherein

- R_1 is selected from the group consisting of hydrogen, alkyl and cycloalkyl;
- Y_1 , Y_2 , Y_3 , Y_4 and Y_5 are independently selected from the group consisting of H, halogen, lower alkyl, O alkyl, O aryl, NH alkyl, NH aryl, N alkyl alkyl, N alkyl aryl, N aryl aryl with the proviso that at least one of Y_1 , Y_2 , Y_3 , Y_4 and Y_5 is aryl or heteroaryl;
- R_4 is selected from the group consisting of alkyl, arylalkyl, cycloalkylalkyl, aminoalkyl, heteroarylalkyl, heterocycloalkylalkyl, CN, COR_5 , CO_2R_5 and CONR_5R_6 ;

R_5 and R_6 are independently selected from the group consisting of H, alkyl, haloalkyl, cycloalkyl, arylalkyl, cycloalkylalkyl, heteroarylalkyl, and heterocycloalkylalkyl;

Z is selected from the group consisting of O, S and NR_8 ;

5 R_8 is selected from the group consisting of H, CN, sulfonamido, OR_7 , alkyl, cycloalkyl, aryl, arylalkyl, heterocyclyl, heteroaryl and heteroarylalkyl; and

R_7 is selected from the group consisting of H, alkyl, arylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, and heteroarylalkyl.

In some preferred embodiments, Y_2 is aryl or heteroaryl. In preferred
10 embodiments, Y_2 is an optionally substituted phenyl, pyridyl, thiazolyl, pyrrolyl, or diazolyl. Preferred substituents include Br, Cl, F, methyl, ethyl, or CF_3 .

According to one embodiment of the instant invention, compounds comprise Formula IA, wherein R_4 is alkyl, arylalkyl, CO_2R_5 or $CONR_5R_6$.

According to another embodiment, the instant invention comprises compounds
15 having formula IA wherein R_1 is alkyl; R_4 is selected from the group consisting of alkyl, arylalkyl, CO_2R_5 , and $CONR_5R_6$; R_5 and R_6 are independently selected from the group consisting of H, alkyl, haloalkyl, cycloalkyl, and arylalkyl; Z is selected from the group consisting of O, S, and NR_8 ; R_8 is selected from the group consisting of H and CN.

20 According to another embodiment, the instant invention comprises compounds of Formula IA wherein R_4 is CO_2R_5 or $CONR_5R_6$ and Z is O.

In other embodiments, the instant invention comprises compounds of formula IA wherein R_4 is selected from the group consisting of alkyl and arylalkyl, and Z is O.

In still further embodiments, the instant invention comprises compounds of
25 formula IA wherein R_1 is CH_3 ; R_4 is CO_2R_5 ; R_5 is alkyl; and Z is O.


The invention further provides a pharmaceutical composition comprising a compound of formula I or IA, as defined above, and a pharmaceutically acceptable carrier. Optionally the pharmaceutical composition may further comprise at least one other anti-cancer agent formulated as a fixed dose.

30 The invention also provides a method for treating a proliferative disease via modulation of the Eg5 motor protein, and/or, inducing apoptosis comprising administering to a mammalian species in need of such treatment an effective amount

of at least one compound of formula I, as defined above. In another embodiment, the invention provides a method for treating a proliferative disease via modulation of the Eg5 motor protein comprising administering to a mammalian species in need of such treatment an effective amount of at least one compound of formula I or Ia, as defined above, in combination (simultaneously or sequentially) with at least one other anti-cancer agent. In a preferred embodiment, the proliferative disease is cancer.

Certain compounds of formula I may generally be prepared according to the following schemes and the knowledge of one skilled in the art. Solvates (e.g., hydrates) of the compounds of formula I and IA are also within the scope of the present invention. Methods of solvation are generally known in the art. Accordingly, the compounds of the instant invention may be in the free or hydrate form, and may be obtained by methods exemplified by the following schemes below.

Scheme I




Compounds of formula I where Z is S may be made in accordance with Scheme I. A ketone or an aldehyde I (e.g., benzaldehyde, where R_2 is phenyl and R_3

is H), is condensed with an acetoacetamide III to give a Knoevenagel product IV as a mixture of isomers. Reaction with S-paramethoxybenzyl thiourea provides the protected dihydropyrimidine thione V. The primary amido group of V is dehydrated to the cyano substituent in VI using a dehydrating agent such as Burgess' reagent

5 (methoxycarbonylsulfamoyl)triethylammonium hydroxide, inner salt. The N3 substituent is introduced by reaction with R_4X where R_4 is alkyl or acyl, and X is a leaving group, or where R_4X is an isocyanate or haloformate. The protecting group is removed by treatment with acid in the presence of water to give compounds of formula I where Z is S.

10




Compounds of formula I where Z is O, NH, or NR_8 are prepared from the reaction of Knoevenagel products IV with O-methyl isourea to provide the O-methyl dihydropyrimidines VIII. The primary amide is converted to a nitrile group using a

15 dehydrating agent such as Burgess' reagent. The N3 substituent is introduced by reaction with R_4X where R_4 is alkyl or acyl, and X is a leaving group, or where R_4X is an isocyanate or haloformate. The methyl ether protecting group is removed by treatment with acid in the presence of water to give compounds of formula I where Z

20 is O. Alternatively, treatment of compounds of formula X with ammonium hydroxide in the presence of ammonium acetate, or cyanamide in ethanol, provides compounds of formula I where Z is NH or NR_8 .

Compounds of formula I may also be prepared using the Bignelli reaction (D. J. Brown in *The Pyrimidines*, Wiley: New York, 1962, 440).

Scheme III
SOLID PHASE SYNTHESIS




Compounds of formula I could be prepared on solid support as outlined in Scheme III. Starting compound XI denotes a resin-bound benzyl alcohol used for solid support synthesis which is prepared from a Merrifield resin denoted as ●, and 2-methoxy-4-hydroxybenzaldehyde, followed by reduction of the aldehyde with reducing agents such as NaBH₄. The benzyl alcohol is converted into the benzyl chloride using agents such as hexachloroethane and triphenylphosphine in THF to form resins of formula XII. The chloride is displaced with thiourea to form the isothiourethane resin XIII. The resulting resin is treated with excess of ketoamides like acetoamide (III, R₁ is CH₃), in the presence of ketones of formula R₂COR₃ or aldehydes of formula R₂CHO to form the resin-bound pyrimidinethiones of formula XIV. The N3 substituent is introduced using R₄X, where X is a leaving group and R₄ is alkyl or acyl, or R₄X is an isocyanate, or haloformate, in the presence of base to form structures of formula XV. The primary amide can be dehydrated to the cyano group using reagents such as Burgess' reagent to form compounds of formula XVI. The products can be cleaved from the resin using a variety of conditions to form compounds of formula I, where Z is determined by the cleavage method employed. Cleavage in the presence of aqueous acid will form compounds of formula I with Z being O, whereas cleavage under anhydrous acid conditions will form compounds of

formula I with Z being S. Alternatively, treatment of resins with structure XVI with ammonium hydroxide in the presence of ammonium acetate will form compounds of formula I with Z being NH, while treatment with cyanamide, provides compounds of formula I with Z being NHCN.

5

Scheme IV




Compounds of formula XVIII may be prepared from a 3-amino-3-alkyl acrylonitrile XVII using the methods illustrated in Scheme IV. Reaction of a compound of formula XVII with aqueous acid, such as hydrochloric acid, followed by treatment with triethyl orthoformate, provides a compound of formula XVIII. Reaction of a compound of formula XVIII with O-methyl isourea in the presence of a base such as triethylamine, provides a pyrimidine of formula XIX. Pyrimidines of formula XIX may be reacted with organometallic species such as a Grignard reagent, R_3MgBr , in a solvent such as ether or tetrahydrofuran, to give a pyrimidine of formula XX, which is a compound of formula IX wherein R_2 is H. In analogy with Scheme II, a compound of formula XX may be converted into a compound of formula XXII, which is a compound of formula I in which R_4 is ethoxycarbonyl and R_2 is H.

Alternatively, compounds of formula I, wherein $\text{Z} = \text{O}$, may be prepared in accordance with Scheme V. Following the procedure of E.H. Hu *et al* (J. Org. Chem., 1998, 63, 3454-3457), a carbonyl compound of formula Ia is condensed with an acylacetamide of formula III in the presence of urea, cuprous chloride and borontrifluoride etherate to give intermediate XXIII. Dehydration with trifluoroacetic anhydride in pyridine affords nitrile XXIV. The N3 substituent is introduced by

reaction with R_4X where R_4 is alkyl or acyl, and X is a leaving group, or where R_4X is an isocyanate or haloformate to give compounds of formula I. Deprotonation is effected with a base such as sodium hydride or LDA in aprotic solvents such as dimethylformamide or tetrahydrofuran.

5


Scheme V



In yet another procedure, compounds of formula I, wherein $Z = O$, may be prepared in accordance with Scheme VI. A carbonyl compound of formula II is condensed with an acylacetamide of formula III in the presence of urea and polyphosphate ester to directly afford nitrile XXIV. Introduction of the N3 substituent follows the procedure as described in Scheme V. Compounds of formula I, where R_4 is aminocarbonyl, may also be obtained by first forming a reactive intermediate such as a nitrophenylcarbamate of formula XXV, which is subsequently reacted with an amine.

15


Scheme VI



In all of the above schema, a 2-acyl acetonitrile derivative, i.e., R_1COCH_2CN , may be substituted for a compound of formula III. schemes below.

5


Scheme VII (Examples 116-117)



Compounds of formula Ia where R₄ is a carbamate may be made in accordance with Scheme VII. Following the procedure of Method 1, an appropriate aldehyde may be condensed with acetoacetamide and urea in the presence of polyphosphoric ester (PPE) to give a dihydropyrimidone product in high yield.

- 5 Reaction of this compound with lithium diisopropylamide and ethyl chloroformate following the procedure of Method 2 provides predominately the N-1 carbamate, which then undergoes a palladium-catalyzed Suzuki coupling reaction with an aryl or heteroaryl boronic acid to give compounds of the present invention as described by Method 3.

Scheme VIII (Examples 118-121)




10

Similarly, compounds of formula Ia where R₄ is a urea may be made in accordance with Scheme VIII. Reaction of a dihydropyrimidone compound obtained following the procedure of Method 1 with sodium hydride and ethyl isocyanate following the procedure of Method 4 provides exclusively the N-1 urea, which then undergoes a palladium-catalyzed Suzuki coupling reaction with an aryl or heteroaryl boronic acid to give compounds of the present invention as described by Method 3.



15

Scheme IX (Examples 122-139, 144-145)




An alternative reaction sequence is described in Scheme IX, wherein a palladium-catalyzed coupling reaction may be performed prior to addition of the exocyclic urea moiety at the N-1 position on the dihydropyrimidone ring. Thus, the dihydropyrimidone product of Method 1 may be reacted with an appropriate aryl or heteroaryl boronic acid in the presence of a palladium catalyst following Method 5 to afford a biaryl product. Subsequent reaction with an isocyanate in the presence of sodium hydride following the procedure of Method 4 affords the compounds of the present invention in good yield. A variety of isocyanate reagents may be prepared from corresponding carboxylic acid precursors by treatment with thionyl chloride and sodium azide using Method 6.

Scheme X (Examples 140-141)



- A variation of the process of Scheme IX involves formation of the boronic acid derivative of the dihydropyrimidinone prepared by Method 1 and is described in Scheme X. This boronic acid derivative may be prepared by reaction of an appropriate aryl halide with bis(pinacolato)diboron in the presence of a palladium catalyst. Subsequent reaction with an appropriate aryl halide in the presence of additional palladium catalyst affords the desired biaryl compound following the procedure of Method 7. Subsequent reaction with an isocyanate in the presence of sodium hydride following the procedure of Method 4 affords the compounds of the present invention in good yield.

Scheme XI (Examples 142-143)



Oxadiazole-substituted compounds of formula IA may be prepared in accordance with Scheme X. Following the procedure of Method 8, an appropriate aryl aldehyde may be condensed with acetoacetamide and urea in the presence of boron trifluoride, followed by dehydration with trifluoroacetic anhydride to give a dihydropyrimidone compound. Reaction of this dihydropyrimidone with an isocyanate and sodium hydride following the procedure of Method 4 affords the N-1 substituted urea, which may be further reacted with lithium hydroxide to provide a carboxylic acid as described by Method 9. Treatment of the carboxylic acid with hydrazine and BOP reagent affords the corresponding acyl hydrazide, which undergoes cyclization with triethyl orthoacetate or an alkyl acid to afford compounds of the present invention as described by Method 10.

As discussed in the background section, Eg5 is a kinesin-like motor protein that facilitates spindle bipolarity during mitosis of the cell cycle. More specifically, the Eg5 protein acts to sort and bundle microtubules of the mitotic spindle during mitosis. Accordingly, Eg5 participates in cell cycle regulation through the spindle checkpoint during the M phase of the cycle. While not wishing to be bound by any theory, it is believed that the compounds of the instant invention act as Eg5 inhibitors. This is theorized because the compounds of the instant invention induce a monopolar astral array of microtubules (the monoastral phenotype) and it has been shown that when Eg5 activity is absent, the monoastral phenotype forms. Regardless of the mechanism of action, the compounds of the instant invention have been shown to cause disruption of the bipolar spindle, spindle checkpoint initiation, mitotic arrest, programmed cell death and tumor cell proliferation inhibition. Furthermore, the compounds of the invention induce a cell cycle arrest in mitosis that is not transient but rather which progresses into programmed cell death. The compounds also exhibit high potency, inducing mitotic arrest and apoptosis in human cells *in vitro* at concentrations in the low or sub μM range. Additionally, in contrast to microtubule agents, the compounds do not disrupt the dynamic instability of microtubules. The instant invention may therefore more specifically target the mitotic spindle of proliferating cells, which may provide for different toxicity profiles than those of existing anti-cancer drugs.

The compounds according to the invention have pharmacological properties; in particular, the compounds of formula I induce mitotic arrest and are believed to be Eg5 inhibitors. The novel compounds of formula I are thus useful in the therapy of a variety of proliferative diseases (including but not limited to diseases associated with the Eg5 motor protein) such as cancer, autoimmune diseases, viral diseases, fungal diseases, neurodegenerative disorders and cardiovascular disease.

More specifically, the compounds of formula I are useful in the treatment of a variety of cancers, including, but not limited to, the following:

- 10 a) carcinoma, including that of the bladder, breast, colon, kidney, liver, lung, including small cell lung cancer, esophagus, gall bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin, including squamous cell carcinoma;
- 15 b) hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkins lymphoma, non-Hodgkins lymphoma, hairy cell lymphoma and Burkett's lymphoma;
- c) hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia;
- 20 d) tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma;
- e) tumors of the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma and schwannomas; and
- 25 f) other tumors, including melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderoma pigmentosum, keratocanthoma, thyroid follicular cancer and Kaposi's sarcoma.

Due to the key role of motor proteins in the regulation of cellular proliferation in general, inhibitors could act as reversible cytostatic agents which may be useful in the treatment of any disease process which features abnormal cellular proliferation, e.g., benign prostatic hyperplasia, familial adenomatosis polyposis, neuro-
30 fibromatosis, atherosclerosis, pulmonary fibrosis, arthritis, psoriasis, glomerulonephritis, restenosis following angioplasty or vascular surgery, hypertrophic scar

formation, inflammatory bowel disease, transplantation rejection, endotoxic shock, and fungal infections.

Compounds of formulae I and Ia induce apoptosis. The apoptotic response is aberrant in a variety of human diseases. Compounds of formula I, as modulators of apoptosis, will be useful in the treatment of cancer (including but not limited to those types mentioned herein above), viral infections (including but not limited to herpesvirus, poxvirus, Epstein-Barr virus, Sindbis virus and adenovirus), prevention of AIDS development in HIV-infected individuals, autoimmune diseases (including but not limited to systemic lupus, erythematosus, autoimmune-mediated glomerulonephritis, rheumatoid arthritis, psoriasis, inflammatory bowel disease, and autoimmune diabetes mellitus), neurodegenerative disorders (including but not limited to Alzheimer's disease, AIDS-related dementia, Parkinson's disease, amyotrophic lateral sclerosis, retinitis pigmentosa, spinal muscular atrophy and cerebellar degeneration), myelodysplastic syndromes, aplastic anemia, ischemic injury associated with myocardial infarctions, stroke and reperfusion injury, arrhythmia, atherosclerosis, toxin-induced or alcohol related liver diseases, hematological diseases (including but not limited to chronic anemia and aplastic anemia), degenerative diseases of the musculoskeletal system (including but not limited to osteoporosis and arthritis) aspirin-sensitive rhinosinusitis, cystic fibrosis, multiple sclerosis, kidney diseases and cancer pain.

Compounds of formula I and Ia may modulate the level of cellular RNA and DNA synthesis. These agents would therefore be useful in the treatment of viral infections (including but not limited to HIV, human papilloma virus, herpesvirus, poxvirus, Epstein-Barr virus, Sindbis virus and adenovirus).

Compounds of formula I and Ia may be useful in the chemoprevention of cancer. Chemoprevention is defined as inhibiting the development of invasive cancer by either blocking the initiating mutagenic event or by blocking the progression of pre-malignant cells that have already suffered an insult or inhibiting tumor relapse.

Compounds of formula I may also be useful in inhibiting tumor angiogenesis and metastasis.

The instant invention may also inhibit other motor proteins, for example, including but not limited to: those human motor proteins that correspond to, Xk1p2,

MKLP1, CHO1, chromokinesins, Nod, Cenp-E, MCAK, members of the BimC family, and members of the Kar3 family. Additionally, compounds used in the methods of the instant invention may also act as inhibitors of other kinesin or kinesin-like proteins and thus be effective in the treatment of diseases associated with other
5 kinesin or kinesin-like proteins.

The compounds of this invention may also be useful in combination (administered together or sequentially) with known anti-cancer treatments such as radiation therapy or with cytostatic or cytotoxic agents, such as for example, but not limited to, DNA interactive agents, such as cisplatin or doxorubicin; topoisomerase II
10 inhibitors, such as etoposide; topoisomerase I inhibitors such as CPT-11 or topotecan; tubulin interacting agents, such as paclitaxel, docetaxel or the epothilones, either naturally occurring or synthetic; hormonal agents, such as tamoxifen; thymidilate synthase inhibitors, such as 5-fluorouracil; and anti-metabolites, such as methotrexate.

15 If formulated as a fixed dose, such combination products employ the compounds of this invention within the dosage range described below and the other pharmaceutically active agent or treatment within its approved dosage range. Compounds of formulae I and Ia may also be administered sequentially with known anticancer or cytotoxic agents when a combination formulation is inappropriate. The
20 invention is not limited in the sequence of administration; compounds of formulae I and Ia may be administered either prior to or after administration of the known anticancer or cytotoxic agent(s).

ASSAYS

25 The pharmacological properties of the compounds of this invention may be confirmed by a number of pharmacological assays. The exemplified pharmacological assays which follow have been carried out with the compounds according to the invention and their salts. The compounds of formulae I and Ia exhibited antiproliferative activity. Preferred compounds exhibit IC_{50} values less than or equal
30 to about 10 μM .

Cell Culture

Cell lines are maintained in RPMI-1640 plus 10% fetal bovine serum.

72-Hour Proliferation Assay

5 Cells were plated at a density of 3,000-6,000 cells/well, depending upon the cell line used, in a 96-well plate. The cultures were grown overnight. Cells were then treated in triplicate with a seven concentration dose-response curve. The maximum concentration of DMSO never exceeded 0.5%. Cells were exposed to compound for 72 hours. Proliferation was measured using XTT or MTS from Promega. The ovarian,
10 breast, prostate, lung, leukemia, and colorectal human cancer cell lines used in this assay included but were not limited to, for example, A2780S, SKBR3, MDA-MB-231, PC3, LX-1, K562, HT-29, WiDr, HCT-15 and HCT116. The compounds of formula I exhibited activity in the 72-hour cell proliferation assay, inhibiting cell proliferation in one or more of the cell lines listed above with at an IC_{50} less than or equal to about 10
15 μM .

Clonogenic Growth Assay

Colony growth inhibition was measured for A2780 ovarian carcinoma cells using a standard clonogenic assay. Briefly, 200 cells/well were seeded into 6-well
20 tissue culture plates (Falcon, Franklin Lakes, NJ) and allowed to attach for 18 hours. Assay medium consisted of RPMI-1640 plus 10% fetal bovine serum. Cells were then treated in duplicate with a six concentration dose-response curve. The maximum concentration of DMSO never exceeded 0.25%. Cells were exposed to compound for 4, 8 or 24 hours. Compound was then removed and the cells were washed with 2
25 volumes of PBS. The normal growth medium was then replaced. Colonies were fed with fresh media every third day. Colony number was scored on day 10-14 using a Optimax imaging station. The compound concentration required to inhibit 50% or 90% of colony formation (IC_{50} or IC_{90} , respectively) was determined by non-linear regression analysis. The coefficient of variance ($SD/mean, n=3$) = 30%. When
30 exposed to cells for 24 hours, the compounds of formulae I and IA exhibited activity in the clonogenicity assay.

Combination Studies - Clonogenic Growth Assays

Combination studies to examine the use of the Eg5 inhibitors of formulae I and Ia in combination with other antineoplastic agents were conducted essentially the same as the standard colony growth assay with the exception of compound treatment.

- 5 In the combination studies, the cells were treated with both a compound of formulae I and Ia and another antineoplastic agent. The compounds were administered simultaneously or sequentially; both the order of sequence and length of treatment (1 to 24 hours) were varied. Data evaluation was based upon the isobologram analysis and the envelope of additivity, using the line of multiplicity which compares the
10 survival fractions of combination treatments with those of single drug treatments.

Cell Cycle Analysis

- The cell cycle profile of cells treated with compounds of formulae I and Ia was monitored by flow cytometry. Briefly, A2780 ovarian carcinoma cells were seeded at
15 a density of 2×10^5 per well in standard 6 well culture plates and permitted to grow for 17 hours. Cells were then exposed to compounds of formulae I and Ia at varying concentrations for 2 to 24 hours. Following exposure, cell populations were harvested, stained with propidium iodide to determine DNA content and also stained with the appropriate immunological reagent for protein biomarkers of mitosis and
20 apoptosis, including, for example, anti-phospho-ThreonineProline, anti-M Phase Phosphoprotein 2 (MMP2), and anti-p85 PARP. The compounds of formulae I and Ia exhibited activity in the cell cycle profile analysis assay, producing significant increases in mitotic and apoptotic fractions of the cell population.

Immunocytochemistry Assays

- A2780 ovarian carcinoma cells or PTK2 kangaroo rat kidney epithelial cells were plated at a density of 200 to 2000 cells per well in 4 chamber glass slides and allowed to attach overnight. Cells were then treated with compounds of formula I and II, separately, at concentrations of 100 nM to 50 μ M for 4 to 30 hours, fixed and
30 permeabilized for subsequent staining. Stain reagents included, for example, propidium iodide, DAPI, rhodamine phalloidin, anti- α tubulin, anti- β tubulin, anti- γ tubulin, and the appropriate fluorescent-tagged secondary antibodies. Cells were

imaged by fluorescent and confocal fluorescent microscopy. The compounds of formulae I and IA inhibited bipolar spindle formation and induced a monoastal array of microtubules.

Further subject matter of the invention also includes pharmaceuticals for use as described above including controlling cancer, inflammation and arthritis, which contain at least one compound of formula I or IA as defined above or at least one of its pharmacologically acceptable acid addition salts, and the use of a compound of the formula I or IA as defined above for the preparation of a pharmaceutical having activity against proliferative diseases as described previously including against cancer, inflammation and/or arthritis.

The following examples and preparations describe the manner and process of making and using the invention and are illustrative rather than limiting. It should be understood that there may be other embodiments which fall within the spirit and scope of the invention as defined by the claims appended hereto.

Example 1

5-Cyano-3,6-dihydro-4-methyl-6-(3-nitrophenyl)-2-thioxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

A. Step 1

A mixture of 6.42 g of acetoacetamide, 8.0 g of 3-nitrobenzaldehyde, 0.61 ml of acetic acid, and 0.21 ml of piperidine in 30 ml of toluene was heated to reflux. A Dean Stark trap was used to azeotrope the water produced. After refluxed for 2 h, the reaction mixture was cooled to room temperature, with a lot of solid appeared, it was treated with a solution of 300 ml of EtOAc and 25 ml MeOH, the solid was then filtered off, rinsed with 15 ml of EtOAc twice to give 3.1 g of desired product in 25% yield.

B. Step 2

A mixture of 200 mg of the compound of Example 1, Step 1, 198 mg of 2-(4-methoxybenzyl)-2-thiopseudourea HCl salt, 84 mg of sodium acetate in 3.6 ml DMF was heated at 85 °C for 15 h, then cooled to room temperature. The resulting reaction mixture was purified by preparative HPLC using a (YMC S5 ODS 20 X 100 mm)

column, the desired fraction was concentrated to dryness. Saturated NaHCO_3 (50 ml) was added and extracted with EtOAc (3 x 50 ml), combined EtOAc extracts were washed with 30 ml of brine, dried with MgSO_4 , filtered and concentrated under vacuum to give 126.1 mg desired product in 36% yield.

5

C. Step 3

A mixture of the compound of Example 1, Step 2 (86.5 mg) and Burgess reagent (150 mg) in 7.0 ml of anhydrous THF was stirred at room temperature for 1 h, concentrated under vacuum, then purified by preparative HPLC using a YMC S5 (10 ODS 20 X 100 mm) column to give 80.8 mg of desired product in 87% yield.

D. Step 4

To a solution of the compound of Example 1, Step 3 (60 mg) and pyridine (0.1 ml) in 0.6 ml of CH_2Cl_2 , 17 μl of ethylchloroformate was added, after stirring for 2.5 h, 15 another 22 μl of ethylchloroformate was added, the reaction mixture was stirred for 2 h, then 0.3 ml of trifluoroacetic acid was added, the resulting mixture was stirred for another 1 h, and concentrated under vacuum; diluted with DMF, MeOH and a little CH_2Cl_2 , filtered, then purified by preparative HPLC using a (YMC S5 ODS 20 X 100 mm) column to give 22.5 mg of product in 42.7% yield. MS $(\text{M}-\text{H})^+ = 345$. HPLC RT 20 = 2.85 min (YMC S5 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.2% phosphoric acid, 4ml/min, monitoring at 220 nm)

Example 2

25 **5-Cyano-3,6-dihydro-4-methyl-6-(3-nitrophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester**

A. Step 1

10.92 g of NaHCO_3 was added portionwise to a solution of 7.83 g of the compound of Example 1, Step 1 and 7.48g of o-methylisourea hydrogen sulfate in DMF (100 ml), there was gas evolved. The reaction mixture was stirred for 2 h, then heated at 65°C 30 overnight, cooled to room temperature, diluted with 800 ml of EtOAc, washed with water (2 x 100 ml) and brine (1 x 100 ml). The organic layer was dried MgSO_4 ,

filtered and concentrated under vacuum. The resulting residue was triturated in EtOAc-CH₂Cl₂-hexane to give 5.48 g of desired product as solid (56%).

B. Step 2

5 A mixture of the compound of Example 2, Step 1 (209 mg) and Burgess reagent (274.5 mg) in CH₂Cl₂ (5 ml) and THF (10 ml) was stirred overnight. The reaction mixture was concentrated under vacuum, diluted with 150 ml of EtOAc, then washed with saturated NaHCO₃ (2 x 30 ml) and brine (1 x 30 ml), dried with MgSO₄, concentrated under vacuum. The resulting residue was purified silica gel
10 chromatography to give 136 mg (69.4%) of desired product.

C. Step 3

1.23 ml of pyridine was added to a solution of 2.075 g of the compound of Example 2, Step 2 in CH₂Cl₂ (30 ml) under argon at 0 °C, then 0.87 ml of ethyl chloroformate
15 was added slowly. The reaction mixture was warmed to room temperature and stirred for 3 h, diluted with a mixture of saturated of NaHCO₃ (50 ml) and brine (50 ml), extracted with EtOAc three times, the combined layers were washed with brine and dried with MgSO₄, filtered and concentrated under vacuum, purified by silica gel
20 chromatography to give 2.57 g (98%) of desired product.

D. Step 4

2.5 ml of TFA was added to a solution of 1.44 g of the compound of Example 2, Step 3 in CH₃CN (25 ml) and H₂O (2.5 ml), the reaction mixture was stirred for 2 h, a lot of white solid appeared. The solid was filtered off, rinsed with CH₃CN (3 x 20 ml)
25 and hexane (2 x 20 ml), dried in air to give 860 mg (62.2%) desired product. The filtrate was concentrated under vacuum, the solid was recrystallized in CH₃CN to give another 320 mg (23.2%) of product. MS (M-H)⁺ = 329. HPLC RT = 2.53 min (YMC S5 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.2% phosphoric acid, 4ml/min, monitoring at 220 nm)
30

Example 3**5-Cyano-3,6-dihydro-4-methyl-6-(3-nitrophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide.****A. Step 1**

5 To a solution of the compound of Example 2, Step 2 (100 mg; 0.37 mmol) and pyridine (0.74 mmol; 18 μ L) in dichloroethane (40 mL) was added 4-nitrophenyl chloroformate (81 mg; 0.40 mmol) and the resulting solution was stirred at room temperature for 1.5 hours. The reaction mixture was diluted with saturated NaHCO_3 (30 mL), extracted with ethyl acetate (3 x 50 mL), dried (MgSO_4) and concentrated in
10 vacuo to afford a white foam. Purification by chromatography (SiO_2 ; 20% EtOAc/hexane) afforded the desired compound as a colorless foam (99 mg; 62%)

B. Step 2

To a solution of the compound of Example 3, Step 1 (12 mg; 27 μ mol) in THF (0.1
15 mL) was added 2M ethylamine in THF solution (15 μ L; 30 mmol) in one portion at room temperature and the resulting yellow solution was stirred 30 minutes. Dilution of the reaction mixture with methanol (1.8 mL) afforded a yellow solid which was collected by suction filtration and purified by preparative HPLC to afford the title compound as a white solid (20 mg; 22%).

20

In contrast to the method of Example 2 above, in this case the 2-methoxy group hydrolyzed during isolation and purification to afford the dihydropyrimidinone ring without the need for treatment with TFA (Example 2, Step 4).

25

Example 4**5-Cyano-3,6-dihydro-4-methyl-6-(3-nitrophenyl)-2-oxo-1-(1-oxobutyl)-(2H)-pyrimidine****A. Step 1**

23.7 μ L of butyryl chloride was added to a solution of 52 mg of the compound of
30 Example 2, Step 2 and 0.15 mL of pyridine in 0.6 mL of anhydrous CH_2Cl_2 , the reaction mixture was stirred for 1h, then 24 μ L of butyryl chloride was added, the

reaction was stirred for 1.5 h, purified by preparative HPLC using a YMC S5 (ODS 20 x 100 mm) column to give 30 mg desired product.

B. Step 2

- 5 A solution of 30 mg of the compound of Example 4, Step 1, 0.2 ml of H₂O and 0.2 ml of TFA in 1.2 ml CH₃CN was stirred for 1.5 h, it was added another 0.1 ml of TFA and stirred for another 2.5 h. The reaction mixture was concentrated under vacuum, and purified by preparative HPLC using a YMC S5 (ODS 20 x 100 mm) column to give 11.8 mg desired product. MS (M-H)⁺ = 327. HPLC RT = 3.06 min (YMC S5
- 10 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.2% phosphoric acid, 4ml/min, monitoring at 220 nm)

Example 5

- enantio* 5-Cyano-3,6-dihydro-4-methyl-6-(3-nitrophenyl)-2-oxo-1-(2H)-
15 pyrimidinecarboxylic acid, 1-ethyl ester (enantiomer A)
- 53 mg of the compound of Example 2, Step 4 was dissolved in absolute EtOH, preparative chiral separation was carried out using a Chiralcel OD-H S5 (4.6 x 250 mm) column, 20 mg of enantiomer A and 27 mg of enantiomer B were obtained. MS (M-H)⁺ = 329. HPLC-Chiral RT = 10.44 min (Chiralcel OD-H, S5, column 4.6 x250
- 20 mm, 10% MeOH / 10% EtOH /Heptane, 1.0 ml /min, monitoring at 220 nm, 94.7% ee)

Example 6

- enantio* 5-Cyano-3,6-dihydro-4-methyl-6-(3-nitrophenyl)-2-oxo-1-(2H)-
25 pyrimidinecarboxylic acid, 1-ethyl ester (enantiomer B)
- MS (M-H)⁺ = 329. HPLC-Chiral RT = 12.92 min (Chiralcel OD-H, S5, column 4.6 x250 mm, 10% MeOH / 10% EtOH /Heptane, 1.0 ml /min, monitoring at 220 nm, 99.64% ee)

30

Example 7

5-Cyano-3,6-dihydro-4-methyl-6-(3-aminophenyl)-2-oxo-1-(2H)-
pyrimidinecarboxylic acid, 1-ethyl ester

A solution of 12 mg of the compound of Example 1, Step 4 in ethanol was treated with 100 mg of tin (II) chloride and heated to reflux under argon for 90 min., the reaction was cooled down and quenched with saturated NaHCO_3 solution and extracted with EtOAc (3 x 50 ml). The combined organic layer was washed with H_2O , dried with Na_2SO_4 and concentrated under vacuum. It was triturated with hexane and ether to give 8 mg of crude product, which was further purified by preparative HPLC to afford 3 mg of desired product as TFA salt. MS $(\text{M}+\text{H})^+ = 301$. HPLC RT = 1.685 min (YMC S5 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.1% of TFA, 4ml/min, monitoring at 220 nm)

10

Example 8

5-Cyano-3,6-dihydro-4-methyl-6-(3-(N,N-dimethyl)aminophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

A solution of 12 mg of the compound of Example 7 in CH_3CN (1 ml) was added paraformaldehyde (40 mg), sodium cyanoborohydride (30 mg) followed by 2 drops of acetic acid. The reaction mixture was stirred at room temperature for 2 h, then quenched with saturated NaHCO_3 solution and extracted with EtOAc three times. The combined organic layer was washed with brine, dried with Na_2SO_4 and concentrated under vacuum. The resulting residue was purified by preparative HPLC to yield 3.2 mg of desired product as TFA salt. MS $(\text{M}+\text{H})^+ = 329$. HPLC RT = 1.76 min (YMC S5 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.1% of TFA, 4ml/min, monitoring at 220 nm)

20

Examples 9 through 15 were prepared using the methods of Example 2 with the substitution of an appropriate benzaldehyde in Step 1.

25

Example 9

5-Cyano-3,6-dihydro-4-methyl-6-(3-trifluoromethylphenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

HPLC-HI 100% at 2.84 min (YMC S5 ODS column 4.6 x 50 mm, 10 - 90% aqueous methanol over 4 minutes containing 0.1% of TFA , 4 ml/min, monitoring at 220 nm). MS: $[\text{M}+\text{H}]^+ = 354$.

30

Example 10

5-Cyano-3,6-dihydro-4-methyl-6-(2,3-dichlorophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

- 5 HPLC-HI 100% at 3.2 min (YMC S5 ODS column 4.6 x 50 mm, 10 - 90% aqueous methanol over 4 minutes containing 0.1% of TFA , 4 ml/min, monitoring at 220 nm).
MS: $[M-H]^- = 352$.

Example 11

- 10 **5-Cyano-3,6-dihydro-4-methyl-6-(3-methoxyphenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester**

HPLC-HI 100% at 2.42 min (YMC S5 ODS column 4.6 x 50 mm, 10 - 90% aqueous methanol over 4 minutes containing 0.1% of TFA , 4 ml/min, monitoring at 220 nm).
MS: $[M+H]^+ = 316$.

Example 12

- 15 **5-Cyano-3,6-dihydro-4-methyl-6-(3,5-dichlorophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester**

HPLC-HI 87% at 3.26 min (YMC S5 ODS column 4.6 x 50 mm, 10 - 90% aqueous methanol over 4 minutes containing 0.1% of TFA , 4 ml/min, monitoring at 220 nm).
20 MS: $[M-H]^- = 352$.

Example 13

5-Cyano-3,6-dihydro-4-methyl-6-(3,4-dichlorophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

- 25 HPLC-HI 100% at 3.197 min (YMC S5 ODS column 4.6 x 50 mm, 10 - 90% aqueous methanol over 4 minutes containing 0.1% of TFA , 4 ml/min, monitoring at 220 nm).
MS: $[M-H]^- = 352$.

Example 14

- 30 **5-Cyano-3,6-dihydro-4-methyl-6-(3-cyanophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester**

HPLC-HI 93% at 2.32 min (YMC S5 ODS column 4.6 x 50 mm, 10 - 90% aqueous methanol over 4 minutes containing 0.1% of TFA, 4 ml/min, monitoring at 220 nm).

MS: $[M+H]^+ = 311$.

5 **Example 15**

5-Cyano-3,6-dihydro-4-methyl-6-(4-methoxyphenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

HPLC-HI 100% at 2.55 min (YMC S5 ODS column 4.6 x 50 mm, 10 - 90% aqueous methanol over 4 minutes containing 0.1% of TFA, 4 ml/min, monitoring at 220 nm).

10 MS: $[M+H]^+ = 316$.

Example 16

5-Cyano-3,6-dihydro-4-methyl-6-(4-methylphenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

15 **A. Step 1**

A cloudy solution of 3-aminocrotononitrile (41 g, 0.5 mol) in Et₂O (500 ml) was added dropwise to the 15% HCl solution (115 ml) at 0°C over 30 min with vigorous stirring, and the reaction mixture was stirred at 0°C for 15 min. The aqueous solution was then separated, extracted with Et₂O (2 x 125 ml), the combined organic phases
20 dried with Na₂SO₄. Triethyl orthoformate (83 ml) in a 500 ml three-neck flask equipped with addition funnel and distillation set was stirred in 60°C-65°C oil bath, the above ether solution was added dropwise such that the rate of addition was equal to the rate of distillation. An additional 83 ml of triethyl orthoformate was added to the reaction when the addition of the ether solution was half complete, the oil bath
25 temperature was slowly raised to 100°C, and the reaction mixture was then stirred for 5 h. Distillation gave 26.6 g (38%) of desired red solid product at 150-155°C/2mm Hg.

B. Step 2

30 To a mixture of O-methylisourea sulfate (9.9 g, 80 mmol), the compound of Example 16, Step 1 (7.4 g, 53 mmol) and ethanol (90 ml) was added Et₃N (11 ml, 80 mmol). The mixture was stirred at room temperature for 15 min, then stirred at 66°C for 3 h,

and concentrated to remove EtOH. EtOAc (80 ml) and H₂O (80 ml) were added, the aqueous layer were separated and extracted with EtOAc (2 x 80 ml), the combined organic layer were dried with Na₂SO₄, concentrated to give brown solid, which was dissolved in EtOAc, filtered through a silica gel pad, washed with EtOAc/heptane (1/1) to remove dark color, and the combined filtrate was concentrated. The solid thus obtained was recrystallized in heptane / EtOAc to give yellow crystal 5.18 g in 65% yield.

C. Step 3.

10 A solution of p-tolylmagnesium bromide in ether (1M, 1 ml, 1 mmol) was added dropwise to a solution of the compound of Example 16, Step 2 (75 mg, 0.5 mmol) in THF (2 ml) at 0°C under argon. The reaction mixture was stirred at the temperature for 1.5 h, another 3 ml of Grignard reagent was added at -78°C, the reaction was slowly warmed to room temperature and stirred for 2 min. Saturated NH₄Cl (5 ml) and H₂O (5 ml) were added, the mixture was extracted with EtOAc (2 x 15 ml), the combined organic layer was dried, concentrated and chromatographed on silica gel to give 45.6 mg of desired product in 91% yield.

D. Step 4

20 To a solution of the compound of Example 16, Step 3 (109 mg, 0.45 mmol) was added pyridine (0.2 ml, 2.5 mmol) in dry CH₂Cl₂ (5 ml) followed by ethyl chloroformate (0.1 ml, 1.05 mmol), and the resulting reaction mixture was stirred at room temperature overnight. MeOH was added, the resulting mixture was stirred for 15 min, concentrated, and chromatographed on silica gel column to give 100 mg desired product as colorless oil (71%).

E. Step 5

A mixture of the compound of Example 15, Step 4 (100 mg, 0.32 mmol), H₂O (0.7 ml), CH₃CN (0.5 ml) and TFA (7 ml) was stirred at room temperature for 2 h. The solution was then concentrated to remove CH₃CN, saturated NaHCO₃ was added to make the mixture basic, the white solid precipitate was filtered, washed with H₂O, and dried to give the desired product (64 mg). The crude product was dried and

recrystallized from EtOH / H₂O to give another 20 mg desired product as white solid. MS (M+H)⁺ = 300. HPLC RT = 3.40 min (YMC S5 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.2% phosphoric acid, 4 ml/min, monitoring at 220 nm)

5

Example 17

5-Cyano-3,6-dihydro-4-methyl-6-cyclohexyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

To a solution of the compound of Example 16, Step 2 (30 mg, 0.2 mmol) in THF (1.2 ml) was added cyclohexylmagnesium chloride (2 M in ether, 1.0 ml, 2 mmol) at -44°C under argon, the reaction was slowly warmed to room temperature, and stirred for 10 min. Saturated NH₄Cl was added, the resulting mixture was extracted several times with EtOAc, the combined organic layer dried, filtered through a silica gel pad, and concentrated to give yellow oil. The oil was dissolved in CH₂Cl₂ (2 ml), then pyridine (80 µl, 0.9 mmol) and ethyl chloroformate (50 µl, 0.5 mmol) were added, the mixture was stirred at room temperature for 30 min, stirred for another 10 min after which H₂O (25 µl) and EtOAc were added, and the mixture dried over Na₂SO₄, filtered through a silica gel pad, and concentrated to give yellow oil. The oil was dissolved in CH₃CN (2 ml), H₂O (0.3 ml) and TFA (0.2 ml) were added, and the mixture stirred at room temperature for 2 h. Saturated NaHCO₃ solution and EtOAc were added, the aqueous layer was separated and extracted with EtOAc, and the combined organic layer was dried over Na₂SO₄, concentrated and chromatographed on silica gel to give 35 mg of desired product as yellow foam (60%). MS (M+H)⁺ = 392. HPLC RT = 3.60 min (YMC S5 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.2% phosphoric acid, 4 ml/min, monitoring at 220 nm)

25

Example 18

5-Cyano-3,6-dihydro-4-methyl-6-phenyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

To a solution of the compound of Example 16, Step 2 (55 mg, 0.37 mmol) in dry THF (2 ml) was added phenylmagnesium bromide (2 M in THF, 2 ml, 4 mmol) dropwise at -78°C under argon. After addition, the reaction was slowly warmed to room

30

temperature and stirred for about 10 min, until starting material disappeared. Saturated NH_4Cl solution and H_2O were added, the mixture was extracted with EtOAc for two times, and the combined organic layer was dried over Na_2SO_4 , concentrated and chromatographed on silica gel to give solid intermediate. The solid
5 was dissolved in CH_2Cl_2 (5 ml), pyridine (0.15 ml, 1.8 mmol) and ethyl chloroformate (0.1 ml, 1 mmol) were added, and the reaction mixture was stirred at room temperature for 0.5 h. The reaction was quenched with 50 μl of H_2O , diluted with 5 ml of EtOAc, the resulting mixture was dried over Na_2SO_4 , filtered through silica gel column to give the intermediate as an oil. The oil was dissolved in CH_3CN (5 ml),
10 H_2O (0.5 ml) and TFA (0.4 ml) were added, the reaction mixture stirred for 1.5 h, and concentrated in vacuo. Saturated NaHCO_3 solution was added to neutralize the mixture, and the precipitate was then filtered and air dried. Recrystallization in EtOAc / heptane to give 70 mg solid product in 66% yield. MS $(\text{M}+\text{H})^+ = 286$. HPLC RT = 1.28 min. (Phenom-Prime S5 C18 4.6 x 30 mm, 10-90 % aqueous methanol over 2
15 minutes containing 0.1 % TFA, 5 ml/min, monitoring at 220 nm)

Examples 19 through 24 were prepared using the method of Example 18 with the substitution of an appropriate arylmagnesiumhalide.

20

Example 19

5-Cyano-3,6-dihydro-4-methyl-6-(2-methylphenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

MS $(\text{M}+\text{H})^+ = 300$. HPLC RT = 1.41 min. (Phenom-Prime S5 C18 4.6 x 30 mm, 10-90 % aqueous methanol over 2 minutes containing 0.1 % TFA, 5 ml/min, monitoring
25 at 220 nm)

Example 20

5-Cyano-3,6-dihydro-4-methyl-6-(3-chlorophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

30 MS $(\text{M}+\text{H})^+ = 320$. HPLC RT = 1.43 min. (Phenom-Prime S5 C18 4.6 x 30 mm, 10-90 % aqueous methanol over 2 minutes containing 0.1 % TFA, 5 ml/min, monitoring at 220 nm)

Example 21

5-Cyano-3,6-dihydro-4-methyl-6-(3-fluorophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

5 MS (M+H)⁺ = 304. HPLC RT = 1.29 min. (Phenom-Prime S5 C18 4.6 x 30 mm, 10-90 % aqueous methanol over 2 minutes containing 0.1 % TFA, 5 ml/min, monitoring at 220 nm)

Example 22

10 **5-Cyano-3,6-dihydro-4-methyl-6-(4-chlorophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester**

MS (M+H)⁺ = 320. HPLC RT = 1.44 min. (Phenom-Prime S5 C18 4.6 x 30 mm, 10-90 % aqueous methanol over 2 minutes containing 0.1 % TFA, 5 ml/min, monitoring at 220 nm)

15

Example 23

5-Cyano-3,6-dihydro-4-methyl-6-(4-fluorophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

MS (M+H)⁺ = 304. HPLC RT = 3.21 min (YMC S5 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.2% phosphoric acid, 4 ml/min, monitoring at 220 nm)

20

Example 24

5-Cyano-3,6-dihydro-4-methyl-6-(2-fluorophenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

25 MS (M+H)⁺ = 304. HPLC RT = 3.05 min (YMC S5 ODS column 4.6 x 50 mm, 10-90% aqueous methanol over 4 minutes containing 0.2% phosphoric acid, 4 ml/min, monitoring at 220 nm)

Example 25**6-(3,5-Bis-trifluoromethylphenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**5 **A. Step 1**

4-(3,5-Bis-trifluoromethylphenyl)-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylic acid amide. A mixture of 3,5-bis(trifluoromethyl)benzaldehyde (2.42 g, 10.0 mmol), acetoacetamide (1.01 g, 10.0 mmol), urea (0.90 g, 15.0 mmol), copper chloride (0.1 g, 1.0 mmol), boron trifluoride etherate (0.09 mL), AcOH (0.04 mL),
10 and THF (20 mL) was heated at 65 °C for 18 h and cooled to room temperature. The resulting precipitates were collected by vacuum filtration, washed with THF, and air dried to give the title compound as an off-white solid (1.33 g, 36%).

B. Step 2

15 **4-(3,5-Bis-trifluoromethylphenyl)-6-methyl-2-oxo-1,2,3,4-tetrahydro-pyrimidine-5-carbonitrile.** To a mixture of compound from Step 1 (1.10 g, 3.0 mmol) and pyridine (12 mL) at 0 °C was added trifluoroacetic anhydride (1.3 mL, 9.0 mmol) slowly over 10 min. The reaction mixture became a clear brown solution and was stirred for 18 h, then poured into water. The resulting off-white precipitates were
20 collected by vacuum filtration, washed with water, and air dried to give the title compound as a light brown solid (1.06 g, 100%).

C. Step 3

6-(3,5-Bis-trifluoromethylphenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester. To a solution of compound from Step 2
25 (60.0 mg, 0.17 mmol) in THF (2 mL) chilled to -78 °C was added LDA (0.15 mL, 0.30 mmol, 2.0 M in hexanes). The resulting mixture was stirred at -78 °C for 15 min, warmed to -25 °C for 15 min, re-cooled to -78 °C and ethyl chloroformate (20 μ L, 0.20 mmol) was added via a syringe. The reaction mixture was stirred at -78 °C for
30 15 min, gradually warmed to room temperature and stirred for 1 h, quenched with MeOH and concentrated in vacuo. The residue was purified by preparative HPLC

(methanol/water); product fractions were combined and concentrated in vacuo to afford the title compound as an off-white solid (28.4 mg, 40%).

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.64 (s, 1 H), 8.19 (s, 1 H), 7.93 (s, 2 H), 6.15 (s, 1 H), 4.17 (m, 2 H), 2.10 (s, 3 H), 1.17 (t, 3 H, *J* = 7.0 Hz). LC/MS (ES+) 444 (M+Na)⁺.

Example 26

6-Butyl-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester

10 A. Step 1:

4-Hexyl-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carbonitrile.

Polyphosphate ester (1.15 g, 3.89 mmol, 0.30 eq) was added to a sealed tube containing a solution of urea (1.17 g, 19.4 mmol), valeraldehyde (1.38 ml, 12.9 mmol) and acetoacetamide (1.31 g, 12.9 mmol) in THF (15 mL) under argon and heated at 15 75°C for three hours. The reaction mixture was cooled to room temperature, polyphosphate ester (8.80 g, 29.8 mmol) was added and the reaction mixture was heated at 85°C for four hours. The reaction mixture was cooled to room temperature and poured on ice. The aqueous solution was neutralized with 1N NaOH and extracted with chloroform (3X100 mL). The combined organic layers were dried 20 (Na₂SO₄), filtered and concentrated under vacuum to give the title compound (0.940 g, 97%): HRMS 192.1143 (M-H)⁺. The product was used in the next step without further purification.

B. Step 2:

25 **6-Butyl-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester.** LDA (0.585 mL, 1.17 mmol, 2.0 M in hexane) was added dropwise to a solution of compound from Step 1 (0.150 g, 0.777 mmol) in THF (5mL) at -78°C. The reaction mixture was stirred at -78°C for twenty minutes and ethyl chloroformate (0.111 mL, 1.17 mmol) was added in one portion. The reaction mixture was allowed 30 to warm to room temperature and was stirred for sixteen hours. The reaction mixture was quenched with saturated ammonium chloride (35 ml) and the solution was extracted with chloroform (3X100 ml). The combined organic layers were dried

(Na₂SO₄), filtered and concentrated under vacuum. The residue was purified by flash chromatography (eluting 2/1 ethyl acetate/hexanes) to afford the title compound (0.110 g, 53%) as a solid.

¹H-NMR (DMSO-d₆, 400 MHz): δ 0.86 (t, 3 H, J = 6.79), 1.21-1.33 (m, 7H), 1.54-1.65 (m, 2H), 2.04 (s, 3H), 4.17-4.23 (m, 2H), 4.68 (t, 1H, J = 6.42), 10.33 (s, 1H). MS 264 (M-H)⁻. HRMS 266.1506 (M+H)⁺.

Example 27

5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid methylamide

A. Step 1

5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid 4-nitrophenyl ester. To a suspension of 6-Methyl-4-(3-nitrophenyl)-2-oxo-1,2,3,4-tetrahydro-pyrimidine-5-carbonitrile (0.43 g, 1.7 mmol, prepared by the method described in Example 26, Step 1, in 77 % yield) in THF (15 mL) at -78 °C was added LDA (1.25 mL, 2.5 mmol, 2.0 M in hexanes). The resulting mixture was stirred at -78 °C for 15 min, warmed to -25 °C for 15 min, recooled to -78 °C and 4-nitrophenyl chloroformate (0.50 g, 2.5 mmol) was added in one portion. The reaction mixture was stirred at -78 °C for 15 min, gradually warmed to room temperature, quenched with water and extracted with ethyl acetate. The organic extract was washed with brine, dried (MgSO₄) and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel using 50% to 70% ethyl acetate-hexane as eluant to give the title compound as a light yellow solid (0.38 g, 53%).

B. Step 2

5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid methylamide. To a solution of compound from Step 1 (12.5 mg, 0.03 mmol) in THF (1 mL) was added methylamine (4.1 uL, 0.033 mmol, 8 M in EtOH). The reaction mixture was stirred for 2 h and concentrated in vacuo. The residue was purified by preparative HPLC (methanol/water). Product fractions were

combined, concentrated in vacuo and lyophilized to give the title compound as a colorless solid (8.2 mg, 87%).

¹H-NMR (MeOH-d₄, 400 MHz): δ 8.22 (m, 1 H), 8.17 (t, 1 H, J = 1.8 Hz), 7.76 (m, 1 H), 7.65 (t, 1 H, J = 7.9 Hz), 6.30 (s, 1 H), 2.82 (s, 3 H), 2.18 (s, 3 H). LC/MS (ES+) 338 (M+Na)⁺.

Example 28

5-Cyano-4-methyl-2-oxo-6-phenyl-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide. To a solution 6-Methyl-2-oxo-4-phenyl-1,2,3,4-tetrahydropyrimidine-5-carbonitrile (100 mg, 0.47 mmole, obtained in 65 % yield by the procedure described in Example 26, Step 1) in 4 mL of tetrahydrofuran under argon at room temperature was added oil free sodium hydride (17 mg, 0.70 mmole). After stirring for ten minutes, ethyl isocyanate (50 mg, 0.70 mmole) was added. The reaction mixture was diluted with ethyl acetate and washed with dilute citric acid, water and saturated brine. The dried (anhydrous magnesium sulfate) organic extract was concentrated *in vacuo* and the product purified by flash chromatography on silica gel, eluting with ethyl acetate/hexanes (3:2) to give 100 mg (75 %) of title compound as a white solid. ¹H-NMR (DMSO-d₆, 400 MHz) δ 10.51 (br s, 1 H); 8.79 (t, J = 5.4 Hz, 1 H); 7.22-7.48 (m, 5 H); 6.09 (s, 1 H); 3.10-3.25 (m, 2 H); 2.08 (s, 3 H); 1.03 (t, J = 7.2 Hz, 3 H). LC/MS (ES+) 285 (M+H)⁺.

Example 29

5-Cyano-6-(3,5-dichlorophenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid 2-methoxy-ethyl ester
Using 3,5-dichlorobenzaldehyde and following the procedure described in Example 2, the title compound was obtained in 3 % overall yield. ¹H-NMR (DMSO-d₆, 500 MHz): δ 10.59 (s, 1 H), 7.66 (t, 1 H, J = 1.6 Hz), 7.31 (d, 2 H, J = 1.6 Hz), 5.90 (s, 1 H), 4.26 (m, 2 H), 3.54 (m, 2 H), 3.25 (s, 3 H), 2.11 (s, 3 H). LC/MS (ES+) 384 (M+H)⁺.

Example 30**5-Cyano-4-methyl-2-oxo-6-(3-trifluoromethylphenyl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 3-trifluoromethylbenzaldehyde and following a modification of the procedure described in Example 2, the title compound was obtained in 3 % overall yield.

¹H-NMR (CDCl₃, 400 MHz): δ 8.68 (s, 1 H), 7.60 (m, 2 H), 7.53 (m, 2 H), 6.37 (s, 1 H), 3.30 (m, 2 H), 2.08 (s, 3 H), 1.17 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 353 (M+H)⁺, 375 (M+Na)⁺.

Example 31**5-Cyano-6-(3,5-dichlorophenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 3,5-dichlorobenzaldehyde and following a modification of the procedure described in Example 2, the title compound was obtained in 5 % overall yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.56 (s, 1 H), 8.76 (t, 1 H, J = 5.3 Hz), 7.63 (t, 1 H, J = 1.8 Hz), 7.27 (d, 2 H, J = 1.8 Hz), 6.10 (s, 1 H), 3.16 (m, 2 H), 2.09 (s, 3 H), 1.03 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 353 (M+H)⁺, 375 (M+Na)⁺.

Example 32**3-Butyryl-4-(3,5-dichlorophenyl)-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carbonitrile**

Using 3,5-dichlorobenzaldehyde and following the procedure described in Example 2, the title compound was obtained in 7 % overall yield.

¹H-NMR (DMSO-d₆, 500 MHz): δ 10.63 (s, 1 H), 7.63 (t, 1 H, J = 1.6 Hz), 7.26 (d, 2 H, J = 1.6 Hz), 6.04 (s, 1 H), 2.95 (m, 1 H), 2.74 (m, 1 H), 1.54 (m, 2 H), 2.11 (s, 3 H), 0.86 (t, 3 H, J = 7.1 Hz).

Example 33**5-Cyano-4-methyl-2-oxo-6-(3-trifluoromethylphenyl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid (2-dimethylamino-ethyl)-amide**

Using 3-trifluoromethylbenzaldehyde and following the procedure described in Example 2, the title compound was obtained in 8 % overall yield.

¹H-NMR (MeOH-d₄, 400 MHz): δ 7.64 (m, 1 H), 7.59 (m, 3 H), 6.25 (s, 1 H), 3.43 (m, 1 H), 3.38 (m, 1 H), 2.56 (t, 1 H, J = 6.2 Hz), 2.32 (s, 6 H), 2.15 (d, 3 H, J = 0.9 Hz); LC/MS (ES+) 396 (M+H)⁺.

5

Example 34**5-Cyano-4-methyl-2-oxo-6-(3-trifluoromethylphenyl)-3,6-dihydro-2H-pyrimidine-1-carbothioic acid S-ethyl ester**

Using 3-trifluoromethylbenzaldehyde and following a modification of the procedure described in Example 2, the title compound was obtained in 3 % overall yield.

10 ¹H-NMR (CDCl₃, 400 MHz): δ 8.21 (s, 1 H), 7.61 (m, 1 H), 7.52 (m, 3 H), 6.24 (s, 1 H), 2.87 (m, 2 H), 2.26 (d, 3 H, J = 0.9 Hz), 1.26 (t, 3 H, J = 7.5 Hz). LC/MS (ES+) 392 (M+Na)⁺.

Example 35

15 **5-Cyano-4-methyl-2-oxo-6-(3-trifluoromethylphenyl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid (2-hydroxy-ethyl)-amide**

Using 3-trifluoromethylbenzaldehyde and following a modification of the procedure described in Example 2, the title compound was obtained in 15 % overall yield.

¹H-NMR (MeOH-d₄, 400 MHz): δ 7.63 (m, 1 H), 7.58 (m, 3 H), 6.25 (s, 1 H), 3.58 (m, 2 H), 3.35 (m, 2 H), 2.15 (s, 3 H). LC/MS (ES+) 369 (M+H)⁺.

20

Example 36**4-(3,5-Dichlorophenyl)-6-methyl-2-oxo-3-(thiophene-2-carbonyl)-1,2,3,4-tetrahydropyrimidine-5-carbonitrile**

25 Using 3,5-dichlorobenzaldehyde and following the procedure described in Example 2, the title compound was obtained in 5 % overall yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.78 (s, 1 H), 7.96 (d, 1 H, J = 4.8 Hz), 7.72 (d, 1 H, J = 3.1 Hz), 7.64 (t, 1 H, J = 1.8 Hz), 7.36 (d, 2 H, J = 1.8 Hz), 7.17 (m, 1 H), 5.92 (s, 1 H), 2.82 (s, 3 H), 2.18 (s, 3 H). LC/MS (ES+) 415 (M+Na)⁺.

30

Example 37**5-Cyano-6-(3,5-difluorophenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Intermediate IX was obtained using 3,5-difluorobenzaldehyde and following the procedure described in Example 2. The title compound was obtained in 28 % yield following the procedure of Example 3.

¹H-NMR (CD₃Cl, 400 MHz): δ 6.77 (m, 2 H), 6.67 (m, 1 H), 6.21 (s, 1 H), 3.27 (m, 2 H), 2.15 (s, 3 H), 1.12 (t, 3H, J = 10.4 Hz). LC/MS (ES+) 321 (M+H)⁺.

Example 38**5-Cyano-6-(3,5-difluorophenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 3,5-difluorobenzaldehyde and following the procedure described in Example 2, the title compound was obtained in 36 % yield for the final two steps.

¹H-NMR (CD₃Cl, 400 MHz): δ 8.26 (s, 1 H), 6.83 (m, 2 H), 6.73 (m, 1 H), 5.80 (s, 1 H), 4.27 (m, 2 H), 2.15 (s, 3 H), 1.26 (t, 3H, J = 10.4 Hz). LC/MS (ES+) 322 (M+H)⁺.

Example 39**4-(3,5-Difluorophenyl)-6-methyl-2-oxo-3-(piperidine-1-carbonyl)-1,2,3,4-tetrahydropyrimidine-5-carbonitrile**

Intermediate IX was obtained using 3,5-difluorobenzaldehyde and following the procedure described in Example 2. The title compound was obtained in 30 % yield following the procedure of Example 3.

¹H-NMR (CD₃OD, 400 MHz): δ 7.01 (m, 3 H), 5.62(s, 1 H), 5.48 (s, 1 H), 3.37 (m, 2 H), 3.24 (m, 2 H), 2.14 (s, 3 H), 1.84 (m, 4), 1.28 (m, 4H). LC/MS (ES+) 361 (M+H)⁺.

Example 40**4-(3,5-Difluorophenyl)-6-methyl-2-oxo-3-(pyrrolidine-1-carbonyl)-1,2,3,4-tetrahydropyrimidine-5-carbonitrile**

Intermediate IX was obtained using 3,5-difluorobenzaldehyde and following the procedure described in Example 2. The title compound was obtained in 18 % yield following the procedure of Example 3.

¹H-NMR (CD₃OD, 400 MHz): δ 8.25 (s, 1 H), 8.24(d, 1 H, J = 7.9 Hz), 7.82 (d, 1 H, J = 7.9 Hz), 7.67 (t, 2 H, J = 7.9 Hz), 5.77 (s, 1H), 2.29-3.30 (m, 4 H), 2.16 (s, 3 H), 1.77 (m, 4). LC/MS (ES+) 355 (M+H)⁺.

5

Example 41**5-Cyano-4-methyl-6-naphthalen-1-yl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using naphthalene-1-carboxaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 44 % yield and the title compound was obtained in 22 % yield.

10

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.57 (s, 1 H), 8.40 (d, 1 H, J = 8.4 Hz), 7.98 (d, 1 H, J = 7.9 Hz), 7.94 (d, 1 H, J = 8.4 Hz), 7.61 (m, 2 H), 7.55 (t, 1 H, J = 7.0 Hz), 7.39 (d, 1 H, J = 7.0 Hz), 6.73 (s, 1 H), 4.04 (m, 2 H), 2.05 (s, 3 H), 1.00 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 336 (M+H)⁺.

15

Example 42**5-Cyano-4-methyl-6-naphthalen-1-yl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using naphthalene-1-carboxaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 44 % yield. The title compound was obtained in 67 % yield following the procedure described in Example 28.

20

¹H-NMR (CDCl₃, 400 MHz): δ 8.83 (t, 1 H, J = 5.3 Hz), 8.43 (d, 1 H, J = 8.4 Hz), 7.96 (s, 1 H), 7.84 (m, 2 H), 7.61 (m, 1 H), 7.52 (t, 1 H, J = 7.0 Hz), 7.42 (t, 1 H, J = 7.9 Hz), 7.36 (d, 1 H, J = 6.2 Hz), 7.14 (s, 1 H), 3.30 (m, 1 H), 3.21 (m, 1 H), 1.92 (s, 3 H), 1.11 (t, 3 H, J = 7.5 Hz). LC/MS (ES+) 335 (M+H)⁺, 357 (M+Na)⁺.

25

Example 43**6-(3-Bromophenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 3-bromobenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 25 % yield and the title compound was obtained in 80 % yield.

30

¹H-NMR (CDCl₃, 400 MHz): δ 7.48 (m, 2 H), 7.27 (m, 2 H), 5.87 (s, 1 H), 4.33 (m, 2 H), 2.22 (s, 3 H), 1.33 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 364 (M+H)⁺.

Example 44

5 **6-(3-Bromophenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 3-bromobenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 25 % yield. The title compound was obtained in 88 % yield following the procedure described in Example 28.

10 ¹H-NMR (CDCl₃, 400 MHz): δ 8.65 (t, 1 H, J = 5.3 Hz), 7.91 (s, 1 H), 7.84 (m, 1 H), 7.38 (m, 1 H), 7.19 (m, 2 H), 6.24 (s, 1 H), 3.24 (m, 2 H), 2.10 (s, 3 H), 1.10 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 363 (M+H)⁺.

Example 45

15 **5-Cyano-4-methyl-2-oxo-6-(2-trifluoromethylphenyl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 2-trifluoromethylbenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 43 % yield and the title compound was obtained in 30 % yield.

20 ¹H-NMR (DMSO-d₆, 400 MHz) δ 10.58 (s, 1 H), 7.77 (m, 2 H), 7.58 (t, 1 H, J = 7.5 Hz), 7.51 (d, 1 H, J = 8.4 Hz), 6.06 (d, 1 H, J = 0.9 Hz), 4.06 (m, 2 H), 2.05 (d, 3 H, J = 0.9 Hz), 1.05 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 354 (M+H)⁺, 376 (M+Na)⁺.

Example 46

25 **5-Cyano-4-methyl-2-oxo-6-(2-trifluoromethylphenyl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 2-trifluoromethylbenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 43 % yield. The title compound was obtained in 26 % yield following the procedure described in Example 28.

30 ¹H-NMR (MeOH-d₄, 400 MHz): δ 9.07 (s, 1 H), 7.69 (d, 1 H, J = 7.9 Hz), 7.64 (d, 1 H, J = 7.9 Hz), 7.54 (d, 1 H, J = 7.9 Hz), 7.49 (t, 1 H, J = 7.9 Hz), 6.48 (s, 1 H), 3.17

(m, 2 H), 2.11 (s, 3 H), 1.06 (t, 3 H, $J = 7.0$ Hz). LC/MS (ES+) 353 (M+H)⁺, 375 (M+Na)⁺.

Example 47

5 **5-Cyano-6-(3-methoxycarbonylphenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using methyl 3-formylbenzoate and following the procedure of Example 25, intermediate XXIV was obtained in 30 % yield and the title compound was obtained in 50 % yield.

10 ¹H-NMR (MeOH-d₄, 400 MHz): δ 8.02 (m, 2 H), 7.60 (m, 1 H), 7.52 (m, 1 H), 5.93 (s, 1 H), 4.28 (m, 2 H), 3.92 (s, 3 H), 2.18 (s, 3 H), 1.29 (t, 3 H, $J = 7.0$ Hz). LC/MS (ES+) 344 (M+H)⁺, 366 (M+Na)⁺.

Example 48

15 **3-(5-Cyano-3-ethylcarbamoyl-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidin-4-yl)-benzoic acid methyl ester**

Using methyl 3-formylbenzoate and following the procedure of Example 25, intermediate XXIV was obtained in 30 % yield. The title compound was obtained in 40 % yield following the procedure described in Example 28.

20 ¹H-NMR (MeOH-d₄, 400 MHz): δ 8.94 (s, 1 H), 7.89 (m, 2 H), 7.48 (d, 1 H, $J = 7.9$ Hz), 7.40 (t, 1 H, $J = 7.9$ Hz), 6.15 (s, 1 H), 3.83 (s, 3 H), 3.17 (m, 2 H), 2.07 (s, 3 H), 1.05 (t, 3 H, $J = 7.0$ Hz). LC/MS (ES+) 343 (M+H)⁺, 365 (M+Na)⁺.

Example 49

25 **6-Butyl-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 1-butanaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 97 % yield. The title compound was obtained in 55 % yield following the procedure described in Example 28.

30 ¹H-NMR (DMSO-d₆, 400 MHz): δ 0.85 (t, 3 H, $J = 6.7$), 1.07 (t, 3 H, $J = 7.2$), 1.23-1.29 (m, 4H), 1.44-1.60 (m, 2H), 2.05 (s, 3H), 3.17-3.32 (m, 2H), 5.04 (t, 1H, $J = 6.0$), 8.70 (t, 1H, $J = 5.5$), 10.30 (s, 1H). MS 263 (M-H)⁻.

Example 50**6-(3,5-Bis-trifluoromethylphenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

- 5 Using 3,5-bis-trifluoromethylbenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 100 % yield. The title compound was obtained in 50 % yield following the procedure described in Example 28.

¹H-NMR (CDCl₃, 400 MHz): δ 8.68 (t, 1 H, J = 5.3 Hz), 7.86 (s, 1 H), 7.77 (s, 2 H), 7.48 (s, 1 H), 6.46 (s, 1 H), 3.31 (m, 2 H), 2.24 (s, 3 H), 1.17 (t, 3 H, J = 7.0 Hz).

10 LC/MS (ES+) 421 (M+H)⁺.

Example 51**6-(2,5-Bis-trifluoromethylphenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

- 15 Using 2,5-bis-trifluoromethylbenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 25 % yield. The title compound was obtained in 38 % yield following the procedure described in Example 28.

¹H-NMR (MeOH-d₄, 400 MHz): δ 7.94 (d, 1 H, J = 8.4 Hz), 7.84 (d, 1 H, J = 8.4 Hz), 7.75 (s, 1 H), 6.50 (s, 1 H), 3.17 (m, 2 H), 2.12 (d, 3 H, J = 0.9 Hz), 1.05 (t, 3 H, J =

20 7.0 Hz). LC/MS (ES+) 421 (M+H)⁺, 443 (M+Na)⁺.

Example 52**6-(2,5-Bis-trifluoromethylphenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

- 25 Using 2,5-bis-trifluoromethylbenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 25 % yield and the title compound was obtained in 55 % yield.

¹H-NMR (CDCl₃, 400 MHz): δ 8.43 (s, 1 H), 7.88 (m, 1 H), 7.76 (m, 2 H), 6.25 (s, 1 H), 4.23 (q, 2 H, J = 7.0 Hz), 2.22 (s, 3 H), 1.19 (t, 3 H, J = 7.0 Hz). LC/MS (ES+)

30 422 (M+H)⁺.

Example 53**5-Cyano-6-(3-ethylcarbamoylphenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Starting with the product of Example 48, the title compound was obtained in 57 %
5 yield by ester hydrolysis (lithium hydroxide in aqueous methanol) and coupling of the
resulting carboxylic acid with ethylamine (water soluble carbodiimide and
hydroxybenzotriazole in acetonitrile).

¹H-NMR (MeOH-d₄, 400 MHz): δ 8.97 (s, 1 H), 7.68 (d, 1 H, J = 1.8 Hz), 7.65 (m, 1
H), 7.38 (m, 2 H), 6.10 (s, 1 H), 3.30 (m, 2 H), 3.13 (m, 2 H), 2.05 (s, 3 H), 1.12 (t, 3
10 H, J = 7.0 Hz), 1.01 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 356 (M+H)⁺.

Example 54**5-Cyano-6-(3-cyano-4-fluorophenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

15 Using 3-cyano-4-fluorobenzaldehyde and following the procedure of Example 25,
intermediate XXIV was obtained in 62 % yield. The title compound was obtained in
51 % yield following the procedure described in Example 28.

¹H-NMR (CDCl₃, 500 MHz) δ 8.66 (s, 1 H), 7.74 (s, 1 H), 7.65 (m, 1 H), 7.57 (m, 1
H), 7.22 (t, 1 H, J = 8.2 Hz), 6.32 (s, 1 H), 3.29 (m, 2 H), 2.22 (s, 3 H), 1.16 (t, 3 H, J
20 = 7.2 Hz). LC/MS (ES+) 328 (M+H)⁺, 350 (M+Na)⁺.

Example 55**5-Cyano-6-(3-cyano-4-fluorophenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

25 Using 3-cyano-4-fluorobenzaldehyde and following the procedure of Example 25,
intermediate XXIV was obtained in 62 % yield and the title compound was obtained
in 30 % yield.

¹H-NMR (MeOH-d₄, 400 MHz): δ 7.71 (m, 2 H), 7.38 (t, 1 H, J = 8.3 Hz), 5.93 (s, 1
H), 4.32 (m, 2 H), 2.20 (d, 3 H, J = 0.9 Hz), 1.32 (t, 3 H, J = 7.0 Hz). LC/MS (ES+)
30 329 (M+H)⁺.

Example 56**5-Cyano-6-(2,2-dimethylpropyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 2,2-dimethylpropionaldehyde and following the procedure of Example 26,
5 intermediate XXIV was obtained in 57 % yield. The title compound was obtained in 54 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δδ 0.92 (s, 9H), 1.06 (t, 3H, J = 7.15Hz), 1.43-1.47 (m, 2H), 2.05 (s, 3H), 3.21 (m, 2H), 5.20 (m, 1H), 8.60 (m, 1H), 10.42 (s, 1H). MS 277 (M-H)⁻.

10

Example 57**5-Cyano-6-cyclopropyl-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using cyclopropanecarboxaldehyde and following the procedure of Example 26,
15 intermediate XXIV was obtained in 83 % yield and the title compound was obtained in 36 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δδ 0.35-0.65 (m, 4H), 1.10-1.21 (m, 1H), 1.23 (t, 3H, J = 7.09 Hz), 2.05 (s, 3H), 4.18-4.21 (m, 2H), 4.33 (d, 1H, J = 8.13 Hz), 10.36 (s, 1H). MS 248 (M-H)⁻.

20

Example 58**5-Cyano-6-cyclopropyl-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using cyclopropanecarboxaldehyde and following the procedure of Example 26,
25 intermediate XXIV was obtained in 83 % yield. The title compound was obtained in 15 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δδ 0.36-0.51 (m, 4H), 1.05-1.09 (m, 4H), 2.06 (s, 3H), 3.20-3.23 (m, 2H), 4.66 (d, 1H, J = 8.30 Hz), 8.74 (m, 1H), 10.36 (s, 1H). MS 247 (M-H)⁻. HRMS 247.1195 (M-H)⁻.

30

Example 59**5-Cyano-6-(2-fluoro-5-trifluoromethylphenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 2-fluoro-5-trifluoromethylbenzaldehyde and following the procedure of

- 5 Example 25, intermediate XXIV was obtained in 41 % yield. The title compound was obtained in 75 % yield following the procedure described in Example 28.

¹H-NMR (CDCl₃, 400 MHz): δ 8.68 (t, 1 H, J = 4.8 Hz), 7.60 (m, 2 H), 7.37 (s, 1 H), 7.22 (t, 1 H, J = 9.2 Hz), 6.41 (s, 1 H), 3.30 (m, 1 H), 3.24 (m, 1 H), 2.18 (d, 3 H, J = 0.9 Hz), 1.14 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 371 (M+H)⁺, 393 (M+Na)⁺.

10

Example 60**5-Cyano-6-(2-fluoro-5-trifluoromethylphenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 2-fluoro-5-trifluoromethylbenzaldehyde and following the procedure of

- 15 Example 25, intermediate XXIV was obtained in 41 % yield and the title compound was obtained in 38 % yield.

¹H-NMR (CDCl₃, 400 MHz): δ 8.27 (s, 1 H), 7.65 (m, 1 H), 7.58 (m, 1 H), 7.25 (t, 1 H, J = 9.2 Hz), 6.03 (s, 1 H), 4.28 (m, 2 H), 2.20 (d, 3 H, J = 0.9 Hz), 1.29 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 372 (M+H)⁺, 394 (M+Na)⁺.

20

Example 61**5-Cyano-6-isopropyl-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using cyclopropanecarboxaldehyde and following the procedure of Example 26,

- 25 intermediate XXIV was obtained in 76 % yield and the title compound was obtained in 23 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 0.88-0.92 (m, 6H), 1.23 (t, 3H, J = 7.10 Hz), 1.90-1.95 (m, 1H), 2.07 (s, 3H), 4.16-4.22 (m, 2H), 4.51 (d, 1H, J = 6.82 Hz), 10.37 (s, 1H). MS: 250 (M-H)⁻.

30

Example 62**5-Cyano-4-methyl-6-naphthalen-2-yl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using naphthalene-2-carboxaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 64 % yield and the title compound was obtained in 39 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.54 (s, 1 H), 7.96 (m, 3 H), 7.78 (s, 1 H), 7.56 (m, 2 H), 7.44 (d, 1 H, J = 8.3 Hz), 5.98 (s, 1 H), 4.18 (q, 2 H, J = 7.0 Hz), 2.12 (s, 3 H), 1.20 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 336 (M+H)⁺.

Example 63**5-Cyano-6-[3-(2-hydroxyethylcarbamoyl)-phenyl]-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Starting with the product of Example 48, the title compound was obtained in 60 % yield by ester hydrolysis (lithium hydroxide in aqueous methanol) and coupling of the resulting carboxylic acid with 2-aminoethanol (water soluble carbodiimide and hydroxybenzotriazole in acetonitrile).

¹H-NMR (CDCl₃, 400 MHz): δ 8.73 (t, 1 H, J = 5.3 Hz), 8.03 (s, 1 H), 7.75 (t, 1 H, J = 1.8 Hz), 7.68 (m, 1 H), 7.51 (d, 1 H, J = 7.5 Hz), 7.39 (t, 1 H, J = 7.5 Hz), 7.00 (t, 1 H, J = 4.8 Hz), 6.29 (s, 1 H), 3.80 (t, 2 H, J = 4.8 Hz), 3.59 (q, 2 H, J = 4.8 Hz), 3.31 (m, 1 H), 3.22 (m, 1 H), 2.18 (s, 3 H), 1.13 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 372 (M+H)⁺, 394 (M+Na)⁺.

Example 64**5-Cyano-6-(2-methoxyphenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 2-methoxybenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 17 % yield and the title compound was obtained in 28 % yield.

¹H-NMR (400 MHz, CDCl₃): δ 1.27 (t, J = 7.0 Hz, 3H), 2.12 (d, J = 0.6Hz, 3H), 3.86 (s, 3H), 4.25 (m, 2H), 6.04 (s, 1H), 6.93 (m, 2H), 7.06 (s, 1H), 7.2 (m, 1H), 7.33 (m, 1H). MS 316 (M+H)⁺.

Example 65

5-Cyano-6-(2-methoxyphenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

Using 2-methoxybenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 17 % yield. Using the procedure of Example 27, the title compound was obtained in 8 % yield.

¹H-NMR (400 MHz, CDCl₃): δ 1.13 (t, J = 7.0 Hz, 3H), 2.1 (s, 3H), 3.25 (m, 1H), 3.33 (m, 1H), 3.86 (s, 3H), 6.47 (s, 1H), 6.62 (s, 1H), 6.9 (m, 2H), 7.2 (dd, J₁ = 1.5 Hz, J₂ = 7.6 Hz, 1H), 7.3 (dt, J₁ = 1.7 Hz, J₂ = 7.2 Hz, 1H), 8.7 (bs, 1H). MS 315 (M+H)⁺.

Example 66

6-(2-Allyloxyphenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester

Using 2-allyloxybenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 33 % yield and the title compound was obtained in 43 % yield.

¹H-NMR (400 MHz, CDCl₃): δ 1.26 (t, J = 7.0 Hz, 3H), 2.1 (s, 3H), 4.25 (m, 2H), 4.6 (m, 2H), 5.30 (d, J = 10.4 Hz, 1H), 5.36 (d, J = 16.0 Hz), 6.0 (s, 1H), 6.1 (m, 1H), 6.9 (m, 3H), 7.25 (m, 2H). MS 342 (M+H)⁺.

Example 67

6-(2-Allyloxyphenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

Using 2-allyloxybenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 33 % yield. Using the procedure of Example 27, the title compound was obtained in 23 % yield.

¹H-NMR (400 MHz, CDCl₃): δ 1.12 (t, J = 7.0 Hz, 3H), 2.1 (s, 3H), 3.2 (m, 1H), 3.3 (m, 1H), 4.6 (m, 2H), 5.35 (m, 2H), 6.1 (m, 1H), 6.38 (s, 1H), 6.50 (s, 1H), 6.9 (m, 2H), 7.25 (m, 2H), 8.7 (bs, 1H). MS 341 (M+H)⁺.

Example 68

6-(2-Bromophenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester

Using 2-bromobenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 50 % yield and the title compound was obtained in 31 % yield.

¹H-NMR (400 MHz, CDCl₃): δ 1.24 (t, J = 7.2 Hz, 3H), 2.16 (s, 3H), 4.24 (q, J = 7.1 Hz, 2H), 6.33 (s, 1H), 7.20 (m, 1H), 7.33 (m, 3H), 7.6 (d, J = 8.0 Hz, 1H). MS 364 (M+H)⁺.

Example 69

6-(2-Bromophenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

Using 2-bromobenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 50 % yield. The title compound was obtained in 46 % yield following the procedure described in Example 28.

¹H-NMR (MeOH-d₄, 500 MHz): δ 7.55 (m, 1 H), 7.30 (m, 2 H), 7.15 (m, 1 H), 6.46 (s, 1 H), 3.14 (m, 2 H), 2.04 (s, 3 H), 1.04 (t, 3 H, J = 7.1 Hz). LC/MS (ES+) 364 (M+H⁺).

Example 70

5-Cyano-6-(3-methoxyphenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

Using 3-methoxybenzaldehyde and following the procedure of Example 25, intermediate XXIV was obtained in 66 % yield. The title compound was obtained in 62 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 500 MHz): δ 10.49 (s, 1 H), 8.79 (t, 1 H, J = 5.5 Hz), 7.32 (t, 1 H, J = 7.7 Hz), 6.92 (m, 1 H), 6.81 (d, 1 H, J = 7.7 Hz), 6.75 (s, 1 H), 6.06 (s, 1 H),

3.74 (s, 3 H), 3.17 (m, 2 H), 2.07 (s, 3 H), 1.04 (t, 3 H, J = 7.1 Hz). LC/MS (ES+) 315 (M+H⁺).

Example 71

5 **6-(2-Benzyloxyphenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 2-benzyloxybenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 47 % yield and the title compound was obtained in 54 % yield.

10 ¹H-NMR (CDCl₃, 400 MHz): δ 1.25 (t, J = 7.0 Hz, 3H), 1.82 (d, J = 0.9 Hz, 3H), 4.21 (m, 2H), 5.04 (dd, J₁ = 10.5 Hz, J₂ = 31.2 Hz, 2H), 5.71 (s, 1H), 5.84 (s, 1H), 6.96 (m, 2H), 7.30 (m, 2H), 7.45 (m, 5H). MS 392 (M+H)⁺.

Example 72

15 **6-Benzyloxymethyl-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 2-benzyloxyacetaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 45 % yield. The title compound was obtained in 30 % yield following the procedure described in Example 28.

20 ¹H-NMR (DMSO-d₆): δ 1.10 (t, 3H, J = 7.33 Hz), 2.07 (s, 3H), 3.22-3.26 (m, 2H), 3.49-3.52 (m, 2H), 4.51 (s, 2H), 5.29 (m, 1H), 7.27-7.34 (m, 5H), 7.90 (br m, 1H), 8.75 (br m, 1H). MS: 327 (M-H)⁻. HRMS 327.1450 (M-H)⁻.

Example 73

25 **6-Benzyloxymethyl-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 2-benzyloxyacetaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 45 % yield and the title compound was obtained in 80 % yield.

30 ¹H-NMR (DMSO-d₆): δ 1.28 (t, 3H, J = 7.13), 2.05 (s, 3H), 3.58-3.70 (m, 2H), 4.29-4.35 (m, 2H), 4.53 (s, 3H), 4.97-4.99 (m, 1H), 7.25-7.35 (m, 5H). MS: 328 (M-H)⁻. 328.1297 HRMS (M-H)⁻.

Example 74**6-(3-Bromo-4-fluorophenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

- 5 Using 3-bromo-4-fluorobenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 99 % yield. The title compound was obtained in 69 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.55 (s, 1 H), 8.75 (t, 1 H, J = 5.3 Hz), 7.55 (m, 1 H), 7.42 (t, 1 H, J = 8.8 Hz), 7.33 (m, 1 H), 6.10 (s, 1 H), 3.16 (m, 2 H), 2.09 (s, 3 H),
10 1.03 (t, 3 H, J = 7.0 Hz). LC/MS (ES+) 381 (M+H⁺).

Example 75**6-(3-Bromo-4-fluorophenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid cyclopropylamide**

- 15 Using 3-bromo-4-fluorobenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 99 % yield. The title compound was obtained in 54 % yield following the procedure described in Example 28.

¹H-NMR (MeOH-d₄, 500 MHz): δ 7.46 (m, 1 H), 7.25 (m, 1 H), 7.14 (t, 1 H, J = 8.8 Hz), 6.04 (s, 1 H), 2.54 (m, 1 H), 2.05 (s, 3 H), 0.63 (m, 2 H), 0.42 (m, 1 H), 0.37 (m,
20 1 H). LC/MS (ES+) 393 (M+H⁺).

Example 76**6-(5-Bromo-2-fluorophenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

- 25 Using 5-bromo-2-fluorobenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 71 % yield. The title compound was obtained in 70 % yield following the procedure described in Example 28.

¹H-NMR (MeOH-d₄, 500 MHz): δ 7.50 (m, 1 H), 7.43 (m, 1 H), 7.10 (m, 1 H), 6.22 (s, 1 H), 3.22 (m, 2 H), 2.11 (d, 3 H, J = 2.2 Hz), 1.11 (m, 3 H).

30

Example 77**6-(5-Bromo-2-fluorophenyl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid cyclopropylamide**

Using 5-bromo-2-fluorobenzaldehyde and following the procedure of Example 26,
5 intermediate XXIV was obtained in 71 % yield. The title compound was obtained in 70 % yield following the procedure described in Example 28.

¹H-NMR (MeOH-d₄, 500 MHz): δ 7.51 (m, 1 H), 7.44 (m, 1 H), 7.11 (m, 1 H), 6.21 (s, 1 H), 2.60 (m, 1 H), 2.10 (d, 3 H, J = 1.6 Hz), 0.70 (m, 2 H), 0.51 (m, 1 H), 0.44 (m, 1 H).

10

Example 78**5-Cyano-6-(3,5-dibromophenyl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 3,5-dibromobenzaldehyde and following the procedure of Example 26,
15 intermediate XXIV was obtained in 46 % yield. The title compound was obtained in 83 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz) δ 10.60 (br s, 1 H); 8.77 (t, J = 5.4 Hz, 1 H); 7.85 (t, J = 1.6 Hz, 1 H); 7.42 (d, J = 1.7 Hz, 2 H); 6.08 (s, 1 H); 3.18-3.26 (m, 2 H); 2.08 (s, 3 H), 1.03 (t, J = 7.1 Hz, 3 H). LC/MS (ES+) 441 (M+H)⁺.

20

Example 79**5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid cyclopropylmethylamide**

Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate
25 XXIV was obtained in 92 % yield. The title compound was obtained in 53 % yield following the procedure described in Example 28.

¹H-NMR (MeOH-d₄, 500 MHz): δ 9.15 (s, 1 H), 8.21 (m, 1 H), 8.17 (d, 1 H, J = 1.6 Hz), 7.76 (d, 1 H, J = 7.7 Hz), 7.66 (t, 1 H, J = 7.7 Hz), 6.29 (s, 1 H), 3.68 (d, 1 H, J = 7.1 Hz), 3.09 (m, 2 H), 2.17 (s, 3 H), 0.98 (m, 1 H), 0.48 (m, 2 H), 0.20 (m, 2 H).

30 LC/MS (ES+) 356 (M+H)⁺.

Example 80**5-Cyano-4-methyl-2-oxo-6-phenethyloxymethyl-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 2-phenethyloxyacetaldehyde and following the procedure of Example 26,
5 intermediate XXIV was obtained in 38 % yield and the title compound was obtained in 18 % yield.

¹H-NMR (DMSO-d₆, 400MHz): δ 1.23 (t, 3H, J = 7.1 Hz), 2.00 (s, 3H), 2.75 (t, 2H, J = 6.7 Hz), 3.59-3.85 (m, 4H), 4.18-4.20 (m, 2H), 4.81 (br m, 1H), 7.19-7.27 (m, 5H), 10.26 (s, 1H). MS 342 (M-H)⁻. HRMS 342.1458 (M-H)⁻.

10

Example 81**5-Cyano-4-methyl-2-oxo-6-phenethyl-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 3-phenylpropionaldehyde and following the procedure of Example 26,
15 intermediate XXIV was obtained in 58 % yield. The title compound was obtained in 42 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δ 1.07 (t, 3H, J = 7.1 Hz), 1.80-2.03 (m, 2H), 2.06 (s, 3H), 2.51-2.75 (m, 2H), 3.19-3.22 (m, 2H), 5.12 (br m, 1H), 7.18-7.30 (m, 5H), 8.70 (br m, 1H), 10.36 (s, 1H). MS 311 (M-H)⁻. HRMS 311.1493 (M-H)⁻.

20

Example 82**5-Cyano-4-methyl-2-oxo-6-thiophen-3-yl-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

25 Using thiophene-3-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 50 % yield and the title compound was obtained in 30 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 1.23 (t, J = 7.1 Hz, 3H), 2.10 (s, 3H), 4.21 (m, 2H), 5.87 (s, 1H), 7.03 (m, 1H), 7.44 (s, 1H), 7.60 (m, 1H), 10.48 (s, 1H). MS 292 (ES⁺):
30 292 (M+H)⁺.

Example 83**5-Cyano-4-methyl-6-(5-methylisoxazol-3-yl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 5-methylisoxazole-3-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 27 % yield and the title compound was obtained in 67 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.52 (s, 1H), 6.20 (s, 1H), 5.95 (s, 1H), 4.21 (q, 2H, J = 7.0 Hz), 2.40 (s, 3H), 2.05 (s, 3H), 1.22 (t, 3H, J = 7.0 Hz). ¹³C-NMR (DMSO-d₆, 400 MHz): δ 170.87, 161.63, 151.86, 150.57, 147.09, 116.44, 99.35, 81.29, 63.08, 49.55, 16.85, 13.68, 11.56. MS (ESI), 291 (M+H)⁺.

Example 84**5-Cyano-6-(3,5-dimethylisoxazol-4-yl)-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 3,5-dimethylisoxazole-4-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 74 % yield and the title compound was obtained in 47 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.56 (s, 1H), 5.82 (s, 1H), 4.15 (q, 2H, J = 7.0 Hz), 2.38 (s, 3H), 2.14 (s, 1H), 2.07 (s, 3H), 1.17 (t, 3H, J = 7.0 Hz). ¹³C-NMR (DMSO-d₆, 400 MHz): δ 167.03, 157.45, 152.17, 147.60, 146.95, 116.44, 113.45, 81.58, 62.96, 48.58, 16.77, 13.64, 10.50, 9.57. MS (ESI), 305 (M+H)⁺.

Example 85**5-Cyano-4-methyl-2-oxo-6-thiophen-2-yl-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using thiophene-2-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 41 % yield and the title compound was obtained in 58 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 1.25 (t, J = 7.1 Hz, 3H), 2.12 (s, 3H), 4.24 (m, 2H), 6.08 (s, 1H), 7.02 (m, 1H), 7.07 (d, J = 3.4 Hz, 1H), 7.56 (m, 1H), 10.58 (s, 1H). MS (ESI) 290 (M-H)⁻.

HRMS (ESI) 290.0598 (M-H)⁻.

Example 86**5-Cyano-4-methyl-6-(3-methylthiophen-2-yl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

- 5 Using 3-methylthiophene-2-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 33 % yield. The title compound was obtained in 17 % yield following the procedure described in Example 28.

¹H NMR (DMSO-d₆, 400 MHz): δδ 1.03 (t, J = 7.2 Hz, 3H), 2.05 (s, 3H), 3.15 (m, 2H), 3.33 (s, 3H), 6.41 (s, 1H), 6.82 (d, J = 5.1 Hz, 1H), 7.35 (d, J = 5.1 Hz, 1H), 8.73 (t, J = 5.6 Hz, 1H), 10.62 (s, 1H).

MS (ESI) 303 (M-H)⁺. HRMS (ESI) 303.0927 (M-H)⁺.

Example 87**5-Cyano-4-methyl-6-(3-methylthiophen-2-yl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

- 15 Using 3-methylthiophene-2-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 33 % yield and the title compound was obtained in 14 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δδ 1.20 (t, J = 7.1 Hz, 3H), 2.05 (s, 3H), 3.33 (s, 3H), 4.17 (q, J = 7.1 Hz, 2H), 6.14 (s, 1H), 6.84 (d, J = 5.1 Hz, 1H), 7.38 (d, J = 5.1 Hz, 1H), 10.61 (s, 1H). MS (ESI) 304 (M-H). HRMS (ESI) 304.0746 (M-H).

Example 88**5-Cyano-4-methyl-6-(2-methylthiazol-4-yl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

- 25 Using 2-methylthiazole-4-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 26 % yield and the title compound was obtained in 16 % yield.

¹H-NMR (CDCl₃, 400 MHz): δ 7.96 (s, 1H), 7.11 (s, 1H), 5.93 (s, 1H), 4.31 (m, 2H), 30 2.67 (s, 3H), 2.16 (s, 3H), 1.32 (t, 3H, J = 6.9 Hz); ¹³C-NMR (CDCl₃, 400 MHz): δ 168.13, 152.34, 151.77, 149.06, 147.71, 116.31, 115.88, 85.39, 64.24, 35.87, 19.17, 17.81, 14.09. MS (ESI) 307 (M+H)⁺.

Example 89**6-(4-Bromothiophen-2-yl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

- 5 Using 4-bromothiophene-2-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 68 % yield. The title compound was obtained in 49 % yield following the procedure described in Example 28. ¹H-NMR (DMSO-d₆, 400 MHz): δδ 1.07 (t, J = 7.1 Hz, 3H), 2.15 (s, 3H), 3.22 (m, 2H), 6.36 (s, 1H), 7.06 (s, 1H), 7.67 (s, 1H), 8.68 (t, J = 5.6 Hz, 1H), 10.64 (s, 1H). MS (ES+) 369 (M+H)⁺.
10 HRMS (ESI): 366.9876 (M-H).

Example 90**6-(4-Bromothiophen-2-yl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

- 15 Using 4-bromothiazole-2-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 68 % yield and the title compound was obtained in 45 % yield. ¹H-NMR (DMSO-d₆, 400 MHz): δδ 1.25 (t, J = 7.1 Hz, 3H), 2.14 (s, 3H), 4.24 (m, 2H), 6.09 (s, 1H), 7.10 (s, 1H), 7.71 (s, 1H), 10.63 (s, 1H). MS (ESI) 368 [M-H]⁻. HRMS (ESI) 367.9697 (M-H).

20

Example 91**5-Cyano-4-methyl-2-oxo-6-[2-(4-trifluoromethylphenyl)-thiazol-4-yl]-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

- Using 2-(4-trifluoromethylphenyl)thiazole-4-carboxaldehyde and following the
25 procedure of Example 26, intermediate XXIV was obtained in 17 % yield and the title compound was obtained in 31 % yield.
¹H-NMR (DMSO-d₆, 400 MHz): δ 10.53 (s, 1H), 8.08 (d, 2H, J = 8.2 Hz), 7.89 (d, 2H, J = 8.1 Hz), 7.78 (s, 1H), 5.96 (s, 1H), 4.20 (m, 2H), 2.07 (s, 3H), 1.22 (t, 3H, J = 7.1 Hz); ¹³C-NMR (DMSO-D₆, 400 MHz): δ 166.18, 154.51, 152.00, 149.60, 142.48,
30 135.79, 130.05, 129.78, 126.45, 125.96, 124.71, 122.30, 117.75, 116.84, 82.69, 62.83, 52.82, 16.87, 13.66. MS (ESI), 437 (M+H)⁺.

Example 92**6-[5-(4-Chlorophenyl)-oxazol-4-yl]-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 5-(4-Chlorophenyl)oxazole-4-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 22 % yield and the title compound was obtained in 16 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.48 (s, 1H), 8.47 (m, 1H), 7.79 (d, 2H, J= 8.6 Hz), 7.64 (d, 2H, J= 8.6 Hz), 6.14 (s, 1H), 4.01 (q, 2H, J= 7.1 Hz), 2.05 (s, 3H), 0.96 (t, 3H, J= 7.1 Hz); ¹³C-NMR (DMSO-D₆, 400 MHz): δ 152.07, 151.98, 149.50, 147.60, 143.20, 133.40, 133.20, 129.01, 127.58, 125.33, 116.82, 81.25, 62.71, 49.80, 16.80, 13.20. MS (ESI), 387 (M+H)⁺.

Example 93**6-(2-Bromothiazol-5-yl)-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester**

Using 2-Bromothiazole-5-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 44 % yield and the title compound was obtained in 41 % yield

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.67 (s, 1H), 7.72 (s, 1H), 6.15 (s, 1H), 4.26 (m, 2H), 2.15 (s, 3H), 1.27 (t, 3H, J= 7.1 Hz); ¹³C-NMR (DMSO-D₆, 400 MHz): δ 152.49, 151.78, 147.11, 141.99, 140.14, 137.93, 116.70, 82.49, 63.91, 50.06, 17.52, 14.18. MS (ESI), 371.0 (M+H)⁺.

Example 94**5-Cyano-4-methyl-2-oxo-6-(4-phenylthiophen-2-yl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 4-bromothiophene-2-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 68 % yield. The 4-bromothiophenyl intermediate was converted to 4-phenylthienyl intermediate by the Suzuki reaction in 91 % yield. The title compound was obtained in 33 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δδ 1.07 (t, J = 7.1 Hz, 3H), 2.17 (s, 3H), 3.23 (m, 2H), 6.42 (s, 1H), 7.30 (t, J = 5.1 Hz, 1H), 7.41-7.43 (m, 3H), 7.66-7.68 (m, 2H), 7.83 (s, 1H), 8.71 (t, J = 5.6 Hz, 1H), 10.61 (s, 1H). MS (ESI) 365 (M-H)⁺. HRMS (ESI) 365.1064(M-H)⁺.

5

Example 95

6-[5-(4-Chlorophenyl)furan-2-yl]-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

Using 5-(4-Chlorophenyl)furan-2-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 16 % yield. The title compound was obtained in 43 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δδ 1.07 (t, J = 7.1 Hz, 3H), 2.12 (s, 3H), 3.22 (m, 2H), 6.27 (s, 1H), 6.46 (d, J = 3.4 Hz, 1H), 6.97 (d, J = 3.4 Hz, 1H), 7.51 (m, 2H), 7.63 (m, 2H), 8.70 (t, J = 5.5 Hz, 1H), 10.62 (s, 1H). MS (ESI) 383 (M-H)⁺. HRMS (ESI) 383.0913 (M-H)⁺.

15

Example 96

5-Cyano-4-methyl-2-oxo-6-(4-trifluoromethylthiazol-5-yl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethyl ester

Using 4-trifluoromethylthiazole-5-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 63 % yield and the title compound was obtained in 52 % yield.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.84 (s, 1H), 9.22 (s, 1H), 6.45 (s, 1H), 4.15 (q, 2H, J = 5.7 Hz), 2.06 (s, 3H), 1.16 (t, 3H, J = 5.7 Hz). ¹³C-NMR (DMSO-D₆, 400 MHz): δ 165.51, 151.90, 149.88, 146.79, 142.70, 138.70, 116.10, 83.68, 63.78, 50.42, 17.44, 13.94; MS (ESI) 361 (M+H)⁺.

25

Example 97

5-Cyano-4-methyl-2-oxo-6-(4-trifluoromethylthiazol-5-yl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

30

Using 4-trifluoromethylthiazole-5-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 63 % yield. The title compound was obtained in 59 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.84 (s, 1H), 9.19 (s, 1H), 8.69 (m, 1H), 6.68 (s, 1H), 3.15 (m, 1H), 2.06 (s, 3H), 1.01 (t, 3H, J = 7.0 Hz); ¹³C-NMR (DMSO-d₆, 400 MHz): δ 155.54, 151.46, 150.25, 149.38, 143.23, 138.70, 115.63, 82.55, 47.93, 34.83, 16.87, 14.34. MS (ESI) 360 (M+H)⁺.

Example 98

5-Cyano-6-[5-(4-methanesulfonylphenyl)-oxazol-4-yl]-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

Using 5-(4-methanesulfonylphenyl)oxazole-4-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 22 % yield. The title compound was obtained in 63 % yield following the procedure described in Example 28.

¹H-NMR (DMF-d₇, 400 MHz): δ 10.60 (s, 1H), 8.96 (t, J = 4.8 Hz, 1H), 8.62 (s, 1H), 8.25 (d, J = 8.3 Hz, 2H), 8.18 (d, J = 8.3 Hz, 2H), 6.67 (s, 1H), 3.37 (s, 3H), 3.25 (m, 2H), 2.21 (s, 3H), 1.07 (t, J = 7.4 Hz, 3H). MS (ESI) 430 (M+H)⁺.

Example 99

6-[5-(4-Chlorophenyl)-oxazol-4-yl]-5-cyano-4-methyl-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

Using 5-(4-chlorophenyl)oxazole-4-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 40 % yield. The title compound was obtained in 58 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.51 (s, 1H), 8.78 (m, 1H), 8.43 (s, 1H), 7.85 (d, 2H, J = 8.5 Hz), 7.62 (d, 2H, J = 8.5 Hz), 6.38 (s, 1H), 3.11 (m, 2H), 2.02 (s, 3H), 0.99 (t, 3H, J = 7.2 Hz). ¹³C-NMR (DMSO-D₆, 400 MHz): δ 152.99, 152.51, 152.11, 149.30, 143.70, 134.17, 129.52, 128.13, 126.40, 117.20, 81.80, 48.09, 35.17, 17.28, 14.95. MS (ESI), 386 (M+H)⁺.

Example 100**5-Cyano-4-methyl-2-oxo-6-[5-(4-trifluoromethylphenyl)-oxazol-4-yl]-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 5-(4-trifluoromethylphenyl)oxazole-4-carboxaldehyde and following the
5 procedure of Example 26, intermediate XXIV was obtained in 31 % yield. The title
compound was obtained in 52 % yield following the procedure described in Example
28.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.54 (s, 1H), 8.78 (m, 1H), 8.50 (s, 1H), 8.06 (d,
2H, J = 8.1 Hz), 7.91 (d, 2H, J = 8.1 Hz), 6.46 (s, 1H), 3.10 (m, 2H), 2.03 (s, 3H), 0.99
10 (t, 3H, J = 7.2 Hz); ¹³C-NMR (DMSO-d₆, 400 MHz): δ 152.53, 152.31, 151.52,
149.03, 142.70, 134.96, 130.69, 126.41, 125.88, 116.62, 81.03, 47.69, 34.65, 16.75,
14.41; MS (ESI), 420.13 (M+H)⁺.

Example 101**5-Cyano-4-methyl-2-oxo-6-(5-phenyloxazol-4-yl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Catalytic reduction (10 % Pd/C, ammonium formate in refluxing acetonitrile) of the
product of Example 99 gave the title compound in 45 % yield.

¹H-NMR (CDCl₃, 400 MHz): δ 8.69 (m, 1H), 7.87 (m, 3H), 7.50 (m, 2H), 7.41 (m,
20 1H), 7.00 (s, 1H), 6.63 (s, 1H), 3.30 (m, 2H), 2.18 (s, 3H), 1.13 (t, 3H, J = 7.2 Hz).
¹³CNMR (CDCl₃, 400 MHz): δ 152.40, 152.13, 150.25, 146.90, 132.60, 129.50,
129.12, 126.79, 116.57, 80.93, 48.25, 35.70, 17.87, 14.72; MS (ESI), 352.16 (M+H)⁺.

Example 102**5-Cyano-4-methyl-6-[5-(4-nitrophenyl)-oxazol-4-yl]-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide**

Using 5-(4-nitrophenyl)oxazole-4-carboxaldehyde and following the procedure of
Example 26, intermediate XXIV was obtained in 36 % yield. The title compound was
obtained in 70 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.56 (s, 1H), 8.79 (m, 1H), 8.56 (s, 1H), 8.38 (m,
30 2H), 8.13 (m, 2H), 6.50 (s, 1H), 3.12 (m, 2H), 2.03 (s, 3H), 0.99 (t, 3H, J = 7.2 Hz).
¹³C-NMR (DMSO-d₆, 400 MHz): δ 153.04, 152.37, 151.51, 149.13, 146.97, 136.16,

132.85, 126.72, 124.19, 116.57, 80.93, 47.87, 34.70, 16.81, 14.45. MS (ESI) 397 (M+H)⁺.

Example 103

5 **5-(4-Chlorophenyl)-2-(5-cyano-3-ethylcarbamoyl-6-methyl-2-oxo-1,2,3,4-tetrahydropyrimidin-4-yl)-furan-3-carboxylic acid ethyl ester**

Using 3-ethoxycarbonyl-5-(4-chlorophenyl)furan-2-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 59 % yield. The title compound was obtained in 42 % yield following the procedure described in Example 10
28.

¹H NMR (DMSO-d₆, 400 MHz): δδ 1.02 (t, J = 7.2 Hz, 3H), 1.34 (t, J = 7.1 Hz, 3H), 2.07 (s, 3H), 3.16 (m, 2H), 4.30 (m, 2H), 6.85 (s, 1H), 7.34 (s, 1H), 7.53 (m, 2H), 7.65 (m, 2H), 8.67 (t, J = 5.5 Hz, 1H), 10.76 (s, 1H). MS (ESI) 455 (M-H)⁻. HRMS (ESI) 455.1114 (M-H)⁻.

15

Example 104

5-Cyano-4-methyl-2-oxo-6-(5-pyridin-3-yloxazol-4-yl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

Using 5-(3-pyridinyl)oxazole-4-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 62 % yield. The title compound was obtained in 41 % yield following the procedure described in Example 28.
20

¹H-NMR (DMSO-d₆, 400 MHz): δδ 10.53 (s, 1H), 9.04 (m, 1H), 8.77 (m, 1H), 8.67 (m, 1H), 8.50 (s, 1H), 8.22 (m, 1H), 7.60 (m, 1H), 6.40 (s, 2H), 3.12 (m, 2H), 2.04 (s, 3H), 1.00 (t, 3H, J = 7.2 Hz). ¹³C-NMR (DMSO-d₆, 400 MHz): □□ 152.76, 152.52, 151.75, 149.97, 149.19, 146.73, 134.73, 133.60, 124.06, 123.45, 116.94, 81.41, 47.81, 34.88, 16.99, 14.63; MS (ESI), 353.18 (M+H)⁺.
25

Example 105

5-Cyano-4-methyl-2-oxo-6-(4-phenylthiazol-5-yl)-3,6-dihydro-2H-pyrimidine-1-carboxylic acid ethylamide

- 5 Using 4-phenylthiazole-5-carboxaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 70 % yield. The title compound was obtained in 27 % yield following the procedure described in Example 28.

¹H-NMR (DMSO-d₆, 400 MHz): δ 10.70 (s, 1H), 9.08 (s, 1H), 8.71 (m, 1H), 7.73 (2H, m), 7.47 (m, 3H), 6.63 (s, 1H), 3.12 (m, 2H), 1.99 (s, 3H), 1.00 (t, 3H, J = 7.2 Hz); ¹³C-NMR (DMSO-d₆, 400 MHz): □□ 152.79, 151.97, 151.74, 150.61, 148.43, 133.87, 116.12, 83.68, 34.77, 16.69, 14.39. MS (ESI) 368 (M+H)⁺.

Example 106

[5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidin-1-yl]-

- 15 **acetic acid tert-butyl ester**

Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 92 % yield and the title compound was obtained in 52 % yield.

¹H-NMR (MeOH-d₄, 400 MHz): δ 8.28 (m, 1 H), 8.22 (t, 1 H, J = 1.8 Hz), 7.76 (m, 1 H), 7.69 (t, 1 H, J = 7.9 Hz), 7.62 (s, 1 H), 5.31 (s, 1 H), 4.23 (d, 1 H, J = 17.6 Hz), 3.44 (d, 1 H, J = 17.6 Hz), 2.16 (s, 3 H), 1.42 (s, 9 H). LC/MS (ES+) 395 (M+Na)⁺.

Example 107

5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid isopropylamide

- 25 Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 92 % yield. The title compound was obtained in 29 % yield following the procedure described in Example 27.

¹H-NMR (MeOH-d₄, 400 MHz): δ 8.92 (s, 1 H), 8.22 (d, 1 H, J = 8.3 Hz), 8.17 (t, 1 H, J = 1.8 Hz), 7.74 (d, 1 H, J = 6.6 Hz), 7.65 (m, 2 H), 6.31 (s, 1 H), 3.89 (m, 1 H), 2.19 (s, 3 H), 1.20 (d, 3 H, J = 6.6 Hz), 1.16 (d, 3 H, J = 6.6 Hz). LC/MS (ES+) 344 (M+H)⁺, 366 (M+Na)⁺.

Example 108**5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid propylamide**

Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate
5 XXIV was obtained in 92 % yield. The title compound was obtained in 22 % yield
following the procedure described in Example 27.

¹H-NMR (MeOH-d₄, 400 MHz) δ 9.09 (t, 1 H, J = 5.3 Hz), 8.22 (m, 1 H), 8.17 (t, 1 H,
J = 1.8 Hz), 7.75 (d, 1 H, J = 7.9 Hz), 7.65 (t, 1 H, J = 7.9 Hz), 6.30 (s, 1 H), 3.33 (m,
2 H), 2.18 (s, 3 H), 1.54 (m, 2 H), 0.91 (t, 3 H, J = 7.5 Hz). LC/MS (ES+) 344
10 (M+H)⁺, 366 (M+Na)⁺.

Example 109**5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid phenylamide**

15 Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate
XXIV was obtained in 92 % yield. The title compound was obtained in 10 % yield
following the procedure described in Example 27.

¹H-NMR (MeOH-d₄, 400 MHz): δ 11.18 (s, 1 H), 8.23 (m, 2 H), 7.81 (d, 1 H, J = 7.9
Hz), 7.65 (m, 1 H), 7.44 (m, 2 H), 7.31 (t, 2 H, J = 8.3 Hz), 7.11 (t, 1 H, J = 8.3 Hz),
20 6.40 (s, 1 H), 2.23 (s, 3 H). LC/MS (ES+) 378 (M+H)⁺, 400 (M+Na)⁺.

Example 110**5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid thiazol-2-ylamide**

25 Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate
XXIV was obtained in 92 % yield. The title compound was obtained in 33 % yield
following the procedure described in Example 27.

¹H-NMR (MeOH-d₄, 400 MHz): δ 8.23 (m, 2 H), 7.80 (d, 1 H, J = 7.5 Hz), 7.64 (t, 1
H, J = 7.5 Hz), 7.42 (d, 1 H, J = 4.0 Hz), 7.03 (d, 1 H, J = 4.0 Hz), 6.36 (s, 1 H), 2.24
30 (d, 3 H, J = 0.9 Hz). LC/MS (ES+) 385 (M+H)⁺, 407 (M+Na)⁺.

Example 111**5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid (2,2,2-trifluoro-ethyl)-amide**

Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 92 % yield. The title compound was obtained in 25 % yield following the procedure described in Example 27.

¹H-NMR (DMSO-d₆, 500 MHz): δ 10.77 (s, 1 H), 9.23 (t, 1 H, J = 6.0 Hz), 8.21 (m, 1 H), 8.09 (s, 1 H), 7.75 (m, 2 H), 6.26 (s, 1 H), 4.01 (m, 2 H), 2.10 (s, 3 H).

10

Example 112**5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid cyclopropylamide**

Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 92 % yield. The title compound was obtained in 19 % yield following the procedure described in Example 27.

¹H-NMR (DMSO-d₆, 500 MHz): δ 10.62 (s, 1 H), 8.58 (s, 1 H), 8.21 (m, 1 H), 8.07 (d, 1 H, J = 1.6 Hz), 7.73 (m, 2 H), 6.22 (s, 1 H), 2.62 (m, 1 H), 2.09 (s, 3 H), 0.64 (m, 2 H), 0.46 (m, 2 H).

20

Example 113**5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid (2-hydroxy-ethyl)-amide**

Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 92 % yield. The title compound was obtained in 23 % yield following the procedure described in Example 27.

¹H-NMR (CDCl₃, 500 MHz): δ 8.93 (s, 1 H), 8.15 (d, 1 H, J = 7.9 Hz), 8.07 (s, 1 H), 7.68 (d, 1 H, J = 7.9 Hz), 7.51 (t, 1 H, J = 7.9 Hz), 6.86 (s, 1 H), 6.30 (s, 1 H), 3.67 (m, 2 H), 3.43 (m, 1 H), 3.36 (m, 1 H), 2.18 (s, 3 H). LC/MS (ES+) 346 (M+H)⁺.

Example 114**3-(4,5-Dihydrooxazol-2-yl)-6-methyl-4-(3-nitrophenyl)-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carbonitrile**

Prepared from the product of Example 113 (Burgess reagent in refluxing tetrahydrofuran) in 71 % yield.

¹H-NMR (MeOH-d₄, 400 MHz): δ 8.10 (m, 2 H), 7.66 (d, 1 H, J = 7.9 Hz), 7.56 (t, 1 H, J = 7.9 Hz), 6.19 (s, 1 H), 4.04 (m, 2 H), 3.50 (m, 1 H), 3.39 (m, 1 H), 2.06 (s, 3 H). LC/MS (ES+) 328 (M+H)⁺.

Example 115**5-Cyano-4-methyl-6-(3-nitrophenyl)-2-oxo-3,6-dihydro-2H-pyrimidine-1-carboxylic acid allylamide**

Using 3-nitrobenzaldehyde and following the procedure of Example 26, intermediate XXIV was obtained in 92 % yield. The title compound was obtained in 36 % yield following the procedure described in Example 27.

¹H-NMR (DMSO-d₆, 500 MHz): δ 10.65 (s, 1 H), 8.92 (t, 1 H, J = 5.5 Hz), 8.21 (m, 1 H), 8.08 (s, 1 H), 7.74 (m, 2 H), 6.24 (s, 1 H), 5.81 (m, 1 H), 5.07 (m, 2 H), 3.78 (m, 2 H), 2.10 (s, 3 H). LC/MS (ES+) 342 (M+H)⁺, 364 (M+Na)⁺.

Example 116**6-(3-Bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine**Method 1

A mixture of 3-bromobenzaldehyde (13.0 g, 70 mmol), acetoacetamide (7.08 g, 70 mmol), urea (5.26 g, 87.5 mmol), polyphosphate ester (9.1 g, 21 mmol) and THF (150 mL) was heated in a sealed tube for 18 h at 75 °C. This mixture was cooled to room temperature; more polyphosphate ester (60.6 g, 140 mmol) was added and heating resumed for another 5 h. The reaction mixture was cooled to room temperature, poured into water (800 mL) and stirred for 18 h. The resulting precipitates were collected by vacuum filtration, washed with water and with ether, and air dried to yield 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine as a light yellow solid (16.25 g; 79%): ¹H NMR (DMSO-d₆, 400 MHz) δ 9.60 (s, 1 H),

7.87 (s, 1 H), 7.55 (m, 1 H), 7.48 (m, 1 H), 7.39 (t, 1 H, $J = 7.9$ Hz), 7.32 (d, 1 H, $J = 7.9$ Hz), 5.14 (s, 1 H), 2.02 (s, 3 H).

6-(3-Bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

Method 2

To a solution of 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine (0.29 g, 1 mmol) in THF (5 mL) chilled to -78°C was added LDA (0.55 mL, 1.1 mmol, 2.0 M in THF). The resulting yellow suspension was stirred for 0.5 h, warmed to -20°C for 0.5 h and recooled to -78°C ; then ethyl chloroformate (0.14 mL, 1.5 mmol) was added via syringe. The reaction mixture was stirred for 20 min, warmed to room temperature and quenched with EtOAc. This mixture was washed with aqueous NaHCO_3 solution and with brine, dried (Na_2SO_4), and concentrated in vacuo. The residue was purified by flash column chromatography (10% to 30% EtOAc/hexanes) to give, after evaporation of solvent, 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester as a colorless solid (0.29 g; 80%): ^1H NMR (CDCl_3 , 400 MHz) δ 7.48 (m, 2 H), 7.28 (m, 2 H), 5.87 (s, 1 H), 4.33 (m, 2 H), 2.22 (s, 3 H), 1.33 (t, 3 H, $J = 7.0$ Hz); LC/MS (ES+) 364 ($\text{M}+\text{H}^+$).

6-(3-Biphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

Method 3

A mixture of 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester (36 mg, 0.1 mmol), phenylboronic acid (24 mg, 0.2 mmol), potassium carbonate (110 mg, 0.8 mmol), EtOH (2 mL), water (1 mL), and toluene (4 mL) was flushed with nitrogen gas several times before palladium tetrakis(triphenylphosphine) (5 mg) was introduced. The reaction mixture was heated at reflux under nitrogen for 1 h, cooled to room temperature, and diluted with EtOAc. This mixture was washed with water, dried (Na_2SO_4), and concentrated in vacuo. The residue was purified by flash column chromatography (50% to 70%

EtOAc/hexanes) to give, after evaporation of solvent, the title compound as a colorless solid (26 mg; 72%): ^1H NMR (MeOH- d_4 , 400 MHz) δ 7.48 (m, 4 H), 7.35 (m, 3 H), 7.23 (m, 2 H), 5.84 (s, 1 H), 4.15 (m, 2 H), 2.04 (s, 3 H), 1.15 (t, 3 H, J = 7.0 Hz); LC/MS (ES+) 362 ($\text{M}+\text{H}^+$).

5

Example 117

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-thienyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester

Following the procedure of Method 3, the title compound was obtained from 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester and 2-thiopheneboronic acid as a light brown solid in 85% yield: ^1H NMR (MeOH- d_4 , 400 MHz) δ 7.52 (m, 2 H), 7.33 (d, 1 H, J = 7.9 Hz), 7.29 (m, 2 H), 7.15 (d, 1 H, J = 7.9 Hz), 6.99 (m, 1 H), 5.81 (s, 1 H), 4.16 (m, 2 H), 2.05 (s, 3 H), 1.17 (t, 3 H, J = 7.0 Hz); LC/MS (ES+) 368 ($\text{M}+\text{H}^+$), 390 ($\text{M}+\text{Na}^+$).

15

Example 118

6-(3-Bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide

20 Method 4

To a solution of 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine (2.92 g, 10 mmol) in THF (30 mL) was added 95% sodium hydride (0.30 g, 12 mmol). After gas evolution ceased, ethyl isocyanate (0.44 mL, 12 mmol) was added via syringe. The reaction mixture was stirred for 0.5 h, quenched with water and extracted twice with EtOAc. The combined extracts were washed with 0.5 N aqueous NaOH, with water and with brine, dried (MgSO_4) and concentrated in vacuo. The residue was purified by flash column chromatography (25% to 35% EtOAc/hexane) to give, after evaporation of solvent, 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide as a colorless solid (2.91 g; 80%): ^1H NMR (CDCl_3 , 400 MHz) δ 8.65 (t, 3 H, J = 5.3 Hz), 7.91 (s, 1 H), 7.39 (m, 1 H), 7.35 (m, 1 H), 7.21 (m, 1 H), 7.15 (d, 1 H, J = 7.9 Hz),

30

6.24 (s, 1 H), 3.25 (m, 2 H), 2.10 (s, 3 H), 1.10 (t, 3 H, $J = 7.0$ Hz); LC/MS (ES+) 363 ($M+H^+$).

5 **6-(3-Biphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide**

Following the procedure of Method 3, the title compound was obtained from 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide and phenylboronic acid as a light yellow solid in 44% yield: 1H NMR ($CDCl_3$, 400 MHz) δ 8.70 (t, 1 H, $J = 5.3$ Hz), 7.55 (m, 4 H), 7.43 (m, 3 H), 7.36 (m, 1 H), 7.29 (d, 1 H, $J = 7.5$ Hz), 7.19 (s, 1 H), 6.39 (s, 1 H), 3.34 (m, 2 H), 2.18 (s, 3 H), 1.17 (t, 3 H, $J = 7.0$ Hz); LC/MS (ES+) 361 ($M+H^+$).

Example 119

15 **6-(3-(4-Chlorophenyl)phenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide**

Following the procedure of Method 3, the title compound was obtained from 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide and 4-chlorophenylboronic acid as a brownish solid in 42% yield: 1H NMR ($MeOH-d_4$, 400 MHz) δ 7.57 (m, 3 H), 7.51 (s, 1 H), 7.45 (m, 3 H), 7.29 (d, 1 H, $J = 7.5$ Hz), 6.21 (s, 1 H), 3.25 (m, 2 H), 2.13 (s, 3 H), 1.12 (t, 3 H, $J = 7.0$ Hz); LC/MS (ES+) 395 ($M+H^+$).

Example 120

25 **5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-thienyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide**

Following the procedure of Method 3, the title compound was obtained from 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide and 2-thiopheneboronic acid as a light yellow solid in 58% yield: 1H NMR ($CDCl_3$, 400 MHz) δ 8.70 (t, 1 H, $J = 5.3$ Hz), 7.54 (m, 2 H), 7.37 (t, 1 H, $J = 7.9$ Hz), 7.30 (m, 2 H), 7.23 (d, 1 H, $J = 7.9$ Hz), 7.12 (s, 1 H), 7.08 (m, 1 H), 6.37 (s, 1 H), 3.32 (m, 2 H), 2.18 (s, 3 H), 1.17 (t, 3 H, $J = 7.0$ Hz); LC/MS (ES+) 367 ($M+H^+$).

Example 121**5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-thienyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide**

- 5 Following the procedure of Method 3, the title compound was obtained from 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide and 3-thiopheneboronic acid as a colorless solid in 14% yield: ^1H NMR (MeOH- d_4 , 400 MHz) δ 7.52 (m, 2 H), 7.46 (s, 1 H), 7.39 (m, 1 H), 7.31 (m, 2 H), 7.11 (d, 1 H, $J = 7.5$ Hz), 6.11 (s, 1 H), 3.17 (m, 2 H), 2.05 (s, 3 H), 1.03 (t, 3 H, $J = 7.0$ Hz); LC/MS $^+$ (ES $^+$) 367 ($\text{M} + \text{H}^+$).
- 10

Example 122**5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-(2H)-pyrimidine**

15 Method 5

- A mixture of 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine (1.62 g, 5.5 mmol), pyridine-4-boronic acid (0.75 g, 6.1 mmol), potassium carbonate (2.28 g, 16.5 mmol) and DMF (10 mL) was flushed with nitrogen gas several times before [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (50 mg) was introduced. The reaction mixture was heated at 95 °C under nitrogen for 64 h, cooled to room temperature and filtered through Celite. The filtrate was diluted with MeOH and stirred with AG 50W-X2 Resin for 10 min. The resin was collected by vacuum filtration, washed with MeOH and then eluted with 2 M NH_3 in MeOH. This NH_3 -MeOH solution was concentrated in vacuo to give 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-(2H)-pyrimidine as a brown solid (1.61 g; 32%); ^1H NMR (DMSO- d_6 , 400 MHz) δ 9.58 (s, 1 H), 8.66 (d, 2 H, $J = 2.2$ Hz), 7.88 (s, 1 H), 7.77 (d, 1 H, $J = 7.9$ Hz), 7.69 (m, 3 H), 7.57 (m, 1 H), 7.41 (d, 1 H, $J = 7.9$ Hz), 5.22 (s, 1 H), 2.03 (d, 3 H, $J = 2.2$ Hz).
- 20
- 25

- 30 **5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide**

Following the procedure of Method 4, the HCl salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-(2H)-pyrimidine and ethyl isocyanate as a light yellow solid in 26% yield: LC/MS (ES+) 362 (M+H⁺).

5

Example 123

2,2,2-Trifluoroethyl isocyanate

Method 6

- 10 A mixture of 3,3,3-trifluoropropionic acid (10.2 g, 80 mmol) and thionyl chloride (5.8 mL, 80 mmol) was heated at reflux for 3 h and cooled to room temperature. Distillation of this mixture afforded 3,3,3-trifluoropropionyl chloride as a colorless liquid (7.12 g; 61%). A mixture of 3,3,3-trifluoropropionyl chloride (6.3 g, 43 mmol), sodium azide (2.8 g, 43 mmol) and toluene (10 mL) was stirred for 18 h at room
- 15 temperature, heated at reflux for 1.5 h and cooled to room temperature to give a solution of 2,2,2-trifluoroethyl isocyanate in toluene (approximately 1.2 M by titration using benzylamine) which was used as such in subsequent reactions.

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-(2,2,2-trifluoro)ethyl amide

- 20 Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-(2H)-pyrimidine and 2,2,2-trifluoroethyl isocyanate as an off-white solid in 45% yield: ¹H NMR (MeOH-d₄, 500 MHz) δδ 9.51 (t, 1 H, J = 6.6 Hz), 8.86 (d, 2 H, J = 6.6 Hz),
- 25 8.30 (d, 2 H, J = 6.6 Hz), 7.93 (d, 1 H, J = 7.7 Hz), 7.90 (s, 1 H), 7.67 (t, 1 H, J = 7.7 Hz), 7.59 (d, 1 H, J = 7.7 Hz), 6.29 (s, 1 H), 3.97 (m, 2 H), 2.18 (s, 3 H); LC/MS (ES+) 416 (M+H⁺).

Example 124

- 30 5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-isopropyl amide

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-(2H)-pyrimidine and isopropyl isocyanate as a colorless solid in 44% yield: ¹H NMR (MeOH-d₄, 400 MHz) δ 8.87 (d, 2 H, J = 7.0 Hz), 8.34 (d, 2 H, J = 7.0 Hz), 7.94 (d, 1 H, J = 7.5 Hz), 7.89 (s, 1 H), 7.68 (t, 1 H, J = 7.5 Hz), 7.58 (d, 1 H, J = 7.5 Hz), 6.27 (s, 1 H), 3.86 (m, 1 H), 2.16 (s, 3 H), 1.18 (d, 3 H, J = 6.6 Hz), 1.12 (d, 3 H, J = 6.6 Hz); LC/MS (ES+) 376 (M+H⁺).

Example 125

10 5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-(2H)-pyrimidine and cyclopropyl isocyanate (prepared from cyclopropanecarbonyl chloride using the procedure of Method 6) as a colorless solid in 42% yield: ¹H NMR (MeOH-d₄, 500 MHz) δ 8.51 (d, 2 H, J = 6.0 Hz), 7.64 (d, 1 H, J = 7.7 Hz), 7.60 (m, 3 H), 7.46 (t, 1 H, J = 7.7 Hz), 7.32 (d, 1 H, J = 7.7 Hz), 6.15 (s, 1 H), 2.54 (m, 1 H), 2.05 (s, 3 H), 0.62 (m, 2 H), 0.43 (m, 1 H), 0.36 (m, 1 H); LC/MS (ES+) 374 (M+H⁺).

20

Example 126

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-(2H)-pyrimidine

Following the procedure of Method 5, 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-(2H)-pyrimidine was obtained from 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine and pyridine-3-boronic acid, 1,3-propanediol cyclic ester as an off-white solid in 76% yield: ¹H NMR (DMSO-d₆, 400 MHz) δ 9.51 (s, 1 H), 8.81 (d, 1 H, J = 2.6 Hz), 8.53 (m, 1 H), 7.99 (m, 1 H), 7.82 (s, 1 H), 7.64 (d, 1 H, J = 7.9 Hz), 7.57 (d, 1 H, J = 1.8 Hz), 7.50 (t, 1 H, J = 7.9 Hz), 7.45 (m, 1 H), 7.30 (d, 1 H, J = 2.6 Hz), 5.15 (s, 1 H), 1.97 (s, 3 H).

30 5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide

Following the procedure of Method 4, the HCl salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-(2H)-pyrimidine and ethyl isocyanate as a light yellow solid in 53% yield: ^1H NMR

(MeOH- d_4 , 400 MHz) δ 9.18 (s, 1 H), 8.92 (m, 1 H), 8.86 (d, 2 H, $J = 5.3$ Hz), 8.19 (m, 1 H), 7.78 (m, 2 H), 7.64 (t, 1 H, $J = 7.5$ Hz), 7.51 (d, 1 H, $J = 7.5$ Hz), 6.24 (s, 1 H), 3.24 (m, 2 H), 2.15 (s, 3 H), 1.11 (t, 3 H, $J = 7.0$ Hz); LC/MS (ES+) 362 ($\text{M} + \text{H}^+$).

Example 127

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-(2,2,2-trifluoro)ethyl amide

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-(2H)-pyrimidine and 2,2,2-trifluoroethyl isocyanate as a colorless solid in 40% yield: ^1H NMR (MeOH- d_4 , 400 MHz) δ 9.51 (s, 1 H), 9.07 (s, 1 H), 8.79 (s, 1 H), 8.71 (d, 1 H, $J = 6.6$ Hz), 8.04 (m, 1 H), 7.76 (d, 1 H, $J = 7.7$ Hz), 7.73 (s, 1 H), 7.62 (m, 1 H), 7.49 (d, 1 H, $J = 7.7$ Hz), 6.27 (s, 1 H), 3.98 (m, 2 H), 2.17 (d, 3 H, $J = 2.7$ Hz); LC/MS (ES+) 416 ($\text{M} + \text{H}^+$).

Example 128

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-isopropyl amide

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-(2H)-pyrimidine and isopropyl isocyanate as a colorless solid in 46% yield: ^1H NMR (MeOH- d_4 , 400 MHz) δ 9.09 (d, 1 H, $J = 1.8$ Hz), 8.79 (d, 1 H, $J = 5.3$ Hz), 8.74 (m, 1 H), 8.05 (m, 1 H), 7.76 (d, 1 H, $J = 7.9$ Hz), 7.73 (d, 1 H, $J = 1.8$ Hz), 7.62 (t, 1 H, $J = 7.9$ Hz), 7.48 (d, 1 H, $J = 7.9$ Hz), 6.24 (s, 1 H), 3.86 (m, 1 H), 2.15 (s, 3 H), 1.18 (d, 3 H, $J = 6.6$ Hz), 1.12 (d, 3 H, $J = 6.6$ Hz); LC/MS (ES+) 376 ($\text{M} + \text{H}^+$).

Example 129

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide

Following the procedure of Method 4, the HCl salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-(2H)-pyrimidine and cyclopropyl isocyanate as an off-white solid in 40% yield: ¹H NMR (MeOH-d₄, 500 MHz) δ 9.08 (s, 1 H), 8.80 (d, 1 H, *J* = 8.2 Hz), 8.77 (d, 1 H, *J* = 5.5 Hz), 8.09 (m, 1 H), 7.70 (d, 1 H, *J* = 7.7 Hz), 7.67 (s, 1 H), 7.55 (t, 1 H, *J* = 7.7 Hz), 7.41 (d, 1 H, *J* = 7.7 Hz) 6.15 (s, 1 H), 2.53 (m, 1 H), 2.06 (s, 3 H), 0.62 (m, 2 H), 0.43 (m, 1 H), 0.36 (m, 1 H); LC/MS (ES+) 374 (M+H⁺).

Example 130

10 **5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclobutyl amide**

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-(2H)-pyrimidine and cyclobutyl isocyanate (prepared from cyclobutanecarbonyl chloride using the procedure of Method 6) as an off-white solid in 88% yield: ¹H NMR (MeOH-d₄, 500 MHz) δ 8.99 (s, 1 H), 8.69 (d, 1 H, *J* = 4.9 Hz), 8.63 (d, 1 H, *J* = 8.2 Hz), 7.95 (m, 1 H), 7.66 (d, 1 H, *J* = 7.7 Hz), 7.62 (s, 1 H), 7.52 (t, 1 H, *J* = 7.7 Hz), 7.38 (d, 1 H, *J* = 7.7 Hz), 6.12 (s, 1 H), 4.09 (m, 1 H), 2.18 (m, 2 H), 2.06 (s, 3 H), 1.82 (m, 2 H), 1.63 (m, 2 H); LC/MS (ES+) 388 (M+H⁺).

20

Example 131

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopentyl amide

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-(2H)-pyrimidine and cyclopentyl isocyanate as an off-white solid in 46% yield: LC/MS (ES+) 402 (M+H⁺).

25 **5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropylmethyl amide**

Example 132

30 **5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropylmethyl amide**

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-(2H)-pyrimidine and cyclopropylmethyl isocyanate (prepared from cyclopropylacetic acid using the procedure of Method 6) as an off-white solid in 62% yield: LC/MS (ES+) (M+H⁺).

Example 133

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-pyridyl)phenyl)-(2H)-pyrimidine

Following the procedure of Method 5, 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-pyridyl)phenyl)-(2H)-pyrimidine was obtained from 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine and 2-tributylstannylpyridine as an off-white solid in 26% yield: ¹H NMR (MeOH-d₄, 500 MHz). 8.71 (d, 1 H, J = 5.0 Hz), 8.25 (m, 1 H), 8.09 (d, 1 H, J = 7.7 Hz), 7.93 (m, 2 H), 7.67 (m, 1 H), 7.63 (d, 1 H, J = 8.2 Hz), 7.57 (d, 1 H, J = 7.7 Hz), 5.29 (s, 1 H), 2.13 (s, 3 H); LC/MS (ES+) 291 (M+H⁺).

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide

Following the procedure of Method 4, the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-pyridyl)phenyl)-(2H)-pyrimidine and ethyl isocyanate as a colorless solid in 25% yield: ¹H NMR (MeOH-d₄, 400 MHz) □ 8.61 (m, 1 H), 7.90 (m, 3 H), 7.82 (d, 1 H, J = 7.9 Hz), 7.51 (m, 1 H), 7.38 (m, 2 H), 6.25 (s, 1 H), 3.25 (m, 2 H), 2.15 (s, 3 H), 1.11 (t, 3 H, J = 7.0 Hz); LC/MS (ES+) 362 (M+H⁺).

Example 134

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-t-butoxycarbonylpyrrole)phenyl)-(2H)-pyrimidine

Following the procedure of Method 5, 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-t-butoxycarbonylpyrrole)phenyl)-(2H)-pyrimidine was obtained from 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine and 1-(t-butoxycarbonyl)pyrrole-2-boronic acid as a light yellow foamy solid in 43% yield: ¹H

NMR (DMSO- d_6 , 400 MHz) δ 9.52 (s, 1 H), 7.84 (s, 1 H), 7.39 (t, 1 H, $J = 7.9$ Hz), 7.34 (m, 1 H), 7.28 (d, 1 H, $J = 7.9$ Hz), 7.23 (d, 1 H, $J = 7.9$ Hz), 7.19 (s, 1 H), 6.28 (m, 1 H), 6.21 (m, 1 H), 5.09 (s, 1 H), 2.00 (s, 3 H), 1.28 (s, 9 H).

5 **5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-pyrrole)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide**

Following the procedure of Method 4 and subsequent treatment with TFA in CH_2Cl_2 for 3 h, the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2- t-butoxycarbonylpyrrole)phenyl)-(2H)-pyrimidine and ethyl isocyanate as a
10 light brown solid in 21% yield: 1H NMR (MeOH- d_4 , 400 MHz) \square 7.49 (m, 2 H), 7.32 (t, 1 H, $J = 7.9$ Hz), 7.06 (d, 1 H, $J = 7.9$ Hz), 6.80 (s, 1 H), 6.43 (s, 1 H), 6.14 (m, 2 H), 3.25 (m, 2 H), 2.14 (s, 3 H), 1.12 (t, 3 H, $J = 7.0$ Hz); LC/MS (ES+) 350 ($M+H^+$).

Example 135

15 **6-(3-Bromo-4-fluorophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine**

Following the procedure of Method 1, 6-(3-bromo-4-fluorophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine was obtained from 3-bromo-4-fluorobenzaldehyde, urea and acetoacetamide as a light yellow solid in 99% yield: 1H
20 NMR (DMSO- d_6 , 400 MHz) δ 9.62 (s, 1 H), 7.86 (s, 1 H), 7.62 (d, 1 H, $J = 6.6$ Hz), 7.44 (t, 1 H, $J = 8.8$ Hz), 7.38 (m, 1 H), 5.18 (s, 1 H), 2.03 (s, 3 H).

5-Cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-(2H)-pyrimidine

25 Following the procedure of Method 5, 5-cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-(2H)-pyrimidine was obtained from 6-(3-bromo-4-fluorophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine and pyridine-3-boronic acid, 1,3-propanediol cyclic ester as a dark brown solid in 44% yield: 1H
30 NMR (DMSO- d_6 , 400 MHz) δ 9.58 (s, 1 H), 8.74 (s, 1 H), 8.62 (m, 1 H), 7.97 (m, 1 H), 7.86 (s, 1 H), 7.53 (m, 2 H), 7.42 (m, 1 H), 7.40 (s, 1 H), 5.23 (s, 1 H), 2.02 (s, 3 H).

5-Cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-
5 (2H)-pyrimidine and ethyl isocyanate as a colorless solid in 52% yield: ^1H NMR (MeOH- d_4 , 400 MHz) δ 8.95 (s, 1 H), 8.74 (d, 1 H, $J = 4.9$ Hz), 8.58 (d, 1 H, $J = 7.7$ Hz), 7.98 (m, 1 H), 7.54 (m, 1 H), 7.43 (m, 1 H), 7.28 (m, 1 H), 6.14 (s, 1 H), 2.06 (s, 3 H), 1.02 (t, 3 H, $J = 7.1$ Hz); LC/MS (ES+) 380 ($\text{M} + \text{H}^+$).

Example 136

5-Cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-(2,2,2-trifluoro)ethyl amide

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-
15 (2H)-pyrimidine and 2,2,2-trifluoroethyl isocyanate as a colorless solid in 31% yield: ^1H NMR (MeOH- d_4 , 500 MHz) \square ^1H NMR (MeOH- d_4 , 400 MHz) δ 9.48 (m, 1 H), 8.99 (s, 1 H), 8.80 (s, 1 H), 8.57 (d, 1 H, $J = 8.2$ Hz), 8.00 (m, 1 H), 7.63 (m, 1 H), 7.53 (m, 1 H), 7.38 (m, 1 H), 6.26 (s, 1 H), 3.97 (m, 2 H), 2.17 (d, 3 H, $J = 1.6$ Hz); LC/MS (ES+) 434 ($\text{M} + \text{H}^+$).

Example 137

5-Cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-
25 (2H)-pyrimidine and cyclopropyl isocyanate as an off-white solid in 39% yield: LC/MS (ES+) 392 ($\text{M} + \text{H}^+$).

Example 138

6-(5-Bromo-2-fluorophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine

Following the procedure of Method 1, 6-(5-bromo-2-fluorophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine was obtained from 5-bromo-2-fluorobenzaldehyde, urea and acetoacetamide as a yellow solid in 72% yield: ^1H NMR (DMSO- d_6 , 400 MHz) δ 9.66 (s, 1 H), 7.81 (s, 1 H), 7.61 (m, 1 H), 7.54 (m, 1 H), 7.27 (m, 1 H), 5.36 (s, 1 H), 2.02 (d, 3 H, $J = 3.1$ Hz).

5-Cyano-3,6-dihydro-6-(2-fluoro-5-(3-pyridyl)phenyl)-4-methyl-2-oxo-(2H)-pyrimidine

Following the procedure of Method 5, 5-cyano-3,6-dihydro-6-(2-fluoro-5-(3-pyridyl)phenyl)-4-methyl-2-oxo-(2H)-pyrimidine was obtained from 6-(5-bromo-2-fluorophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine and pyridine-3-boronic acid, 1,3-propanediol cyclic ester as an off-white solid in 59% yield: ^1H NMR (DMSO- d_6 , 400 MHz) δ 9.62 (s, 1 H), 8.86 (d, 1 H, $J = 1.8$ Hz), 8.58 (m, 1 H), 8.04 (m, 1 H), 7.81 (s, 1 H), 7.77 (m, 1 H), 7.69 (m, 1 H), 7.50 (m, 1 H), 7.38 (m, 1 H), 5.41 (s, 1 H), 2.01 (d, 3 H, $J = 0.9$ Hz).

5-Cyano-3,6-dihydro-6-(2-fluoro-5-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide

Following the procedure of Method 4, the HCl salt of the title compound was obtained from 5-cyano-3,6-dihydro-6-(2-fluoro-5-(3-pyridyl)phenyl)-4-methyl-2-oxo-(2H)-pyrimidine and ethyl isocyanate as a slightly yellow solid in 34% yield: ^1H NMR (MeOH- d_4 , 500 MHz) δ 9.20 (s, 1 H), 8.91 (d, 1 H, $J = 8.2$ Hz), 8.86 (d, 1 H, $J = 4.4$ Hz), 8.19 (m, 1 H), 7.84 (m, 2 H), 7.40 (m, 1 H), 6.29 (s, 1 H), 3.20 (m, 2 H), 2.12 (d, 3 H, $J = 2.2$ Hz), 1.09 (m, 3 H); LC/MS (ES+) 380 ($\text{M} + \text{H}^+$).

Example 139

5-Cyano-3,6-dihydro-6-(2-fluoro-5-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide

Following the procedure of Method 4, the HCl salt of the title compound was obtained from 5-cyano-3,6-dihydro-6-(2-fluoro-5-(3-pyridyl)phenyl)-4-methyl-2-oxo-(2H)-pyrimidine and cyclopropyl isocyanate as a light yellow solid in 32% yield: ^1H

NMR (MeOH-d₄, 500 MHz) δ 9.19 (s, 1 H), 8.91 (d, 1 H, $J = 8.2$ Hz), 8.85 (d, 1 H, $J = 5.5$ Hz), 8.17 (m, 1 H), 7.84 (m, 2 H), 7.40 (m, 1 H), 6.28 (s, 1 H), 2.58 (m, 1 H), 2.11 (s, 3 H), 0.69 (m, 2 H), 0.51 (m, 1 H), 0.42 (m, 1 H); LC/MS (ES+) 392 (M+H⁺).

5

Example 140**5-Cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-3-pyridyl)phenyl)-2-oxo-(2H)-pyrimidine**Method 7

- 10 A mixture of 6-(3-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine (1.46 g, 5 mmol), bis(pinacolato)diboron (1.40 g, 5.5 mmol), potassium acetate (1.47 g, 16.5 mmol) and DMF (5 mL) was flushed with nitrogen gas several times before [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (50 mg) was introduced. The reaction mixture was heated at 95 °C under nitrogen for 18 h, cooled to room temperature and filtered through Celite. The filtrate was diluted with EtOAc, washed with water and with brine, dried (Na₂SO₄) and concentrated to give a dark viscous oil. This material was re-dissolved in DMF (5 mL) and reacted under nitrogen with 5-bromo-2-methylpyridine (0.66 g, 3.8 mmol), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium (50 mg) and potassium carbonate (2.07 g, 15 mmol) at 95 °C for 18 h. The reaction mixture was cooled to room temperature, diluted with MeOH and stirred with AG 50W-X2 Resin for 5 min. The resin was collected by vacuum filtration, washed with MeOH and then eluted with 2 M NH₃ in MeOH. This NH₃-MeOH solution was concentrated in vacuo to give 5-cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-3-pyridyl)phenyl)-2-oxo-(2H)-pyrimidine as a brown solid (0.69 g; 45%): ¹H NMR (DMSO-d₆, 400 MHz) δ 9.55 (s, 1 H), 8.72 (s, 1 H), 7.92 (d, 1 H, $J = 6.2$ Hz), 7.85 (s, 1 H), 7.64 (d, 1 H, $J = 7.5$ Hz), 7.57 (s, 1 H), 7.51 (t, 1 H, $J = 7.5$ Hz), 7.33 (m, 2 H), 5.18 (s, 1 H), 2.49 (s, 3 H), 2.01 (s, 3 H).

- 30 **5-Cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-3-pyridyl)phenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide**

Following the procedure of Method 4, the HCl salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-3-pyridyl)phenyl)-2-oxo-(2H)-pyrimidine and ethyl isocyanate as a slightly yellow solid in 33% yield: ^1H NMR (MeOH- d_4 , 500 MHz) δ 8.99 (d, 1 H, $J = 2.2$ Hz), 8.75 (m, 1 H), 8.01 (d, 1 H, $J = 8.2$ Hz), 7.76 (d, 1 H, $J = 7.7$ Hz), 7.73 (s, 1 H), 7.62 (t, 1 H, $J = 7.7$ Hz), 7.49 (d, 1 H, $J = 7.7$ Hz), 6.23 (s, 1 H), 3.25 (m, 2 H), 2.84 (s, 3 H), 2.15 (s, 3 H), 1.12 (t, 3 H, $J = 7.1$ Hz); LC/MS (ES+) 376 ($\text{M} + \text{H}^+$).

Example 141

10 5-Cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-3-pyridyl)phenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide

Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-3-pyridyl)phenyl)-2-oxo-(2H)-pyrimidine and cyclopropyl isocyanate as a colorless solid in 17% yield: ^1H NMR (MeOH- d_4 , 400 MHz) δ 8.95 (s, 1 H), 8.66 (d, 1 H, $J = 8.3$ Hz), 7.93 (d, 1 H, $J = 8.3$ Hz), 7.75 (d, 1 H, $J = 7.9$ Hz), 7.72 (s, 1 H), 7.61 (t, 1 H, $J = 7.9$ Hz), 7.47 (d, 1 H, $J = 7.9$ Hz), 6.23 (s, 1 H), 2.81 (s, 3 H), 2.63 (m, 1 H), 2.15 (s, 3 H), 0.71 (m, 2 H), 0.51 (m, 1 H), 0.46 (m, 1 H); LC/MS (ES+) 388 ($\text{M} + \text{H}^+$).

20 Example 142

6-(3-Carbomethoxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine

Method 8

- 25 A mixture of 3-carbomethoxybenzaldehyde (4.15 g, 25.3 mmol), acetoacetamide (2.56 g, 25.3 mmol), urea (2.28 g, 38.0 mmol), boron trifluoride etherate (0.25 mL), copper chloride (0.25 g, 2.5 mmol), acetic acid (0.1 mL) and THF (50 mL) was heated at 65 °C for 18 h and cooled to room temperature. The resulting precipitates were collected by vacuum filtration, washed with THF and air dried to give 7.48 g of an off-white solid.
- 30 To a mixture of this solid and pyridine (100 mL) at 0 °C was added trifluoroacetic anhydride (10.6 mL, 7.5 mmol) slowly via syringe. The resulting brown solution was stirred for 2.5 h and poured into water to result in precipitation. The precipitates were

collected by vacuum filtration, washed with water and with CH_2Cl_2 , and air dried to give 6-(3-carbomethoxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine as a light tan solid (4.57 g, 67%): ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ 9.62 (s, 1 H), 7.94 (m, 2 H), 7.90 (s, 1 H), 7.58 (m, 2 H), 5.24 (s, 1 H), 3.87 (s, 3 H), 2.02 (s, 3 H).

5

6-(3-Carbomethoxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide

Following the procedure of Method 4, 6-(3-carbomethoxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide was obtained from 6-(3-carbomethoxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine and ethyl isocyanate as a colorless solid in 52% yield: ^1H NMR ($\text{MeOH}-d_4$, 400 MHz) δ 8.94 (s, 1 H), 7.89 (m, 2 H), 7.48 (d, 1 H, $J = 7.9$ Hz), 7.40 (t, 1 H, $J = 7.9$ Hz), 6.15 (s, 1 H), 3.83 (s, 3 H), 3.17 (m, 2 H), 2.07 (s, 3 H), 1.05 (t, 3 H, $J = 7.0$ Hz); LC/MS (ES+) 343 ($\text{M}+\text{H}^+$), 365 ($\text{M}+\text{Na}^+$).

15

6-(3-Carboxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide

Method 9

To a mixture of 6-(3-carbomethoxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide (0.34 g, 1.0 mmol), MeOH (9 mL) and water (3 mL) at 0 °C was added lithium hydroxide (0.84g, 20.0 mmol). The reaction mixture was stirred for 1 h, acidified to pH 2 with 1 N HCl and extracted with EtOAc. The extract was dried (MgSO_4) and concentrated in vacuo to give 6-(3-carboxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide as a colorless solid (0.30 g, 92%): ^1H NMR ($\text{MeOH}-d_4$, 400 MHz) δ 8.98 (s, 1 H), 7.89 (m, 2 H), 7.46 (d, 1 H, $J = 7.5$ Hz), 7.42 (t, 1 H, $J = 7.5$ Hz), 6.12 (s, 1 H), 3.16 (m, 2 H), 2.06 (s, 3 H), 1.03 (t, 3 H, $J = 7.0$ Hz).

5-Cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-5-oxa-2,3-diazole)phenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide

Method 10

A mixture of 6-(3-carboxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide (0.03 g, 0.1 mmol), BOP reagent (0.07 g, 0.15 mmol) diisopropylethylamine (0.03 mL, 0.15 mmol), hydrazine (5 μ L, 0.15 mmol) and THF (1 mL) was stirred for 1 h and concentrated in vacuo. The residue was heated with triethyl orthoacetate (1 mL) at 80 °C for 18 h to give a light red solution. This solution was concentrated in vacuo; purification of the residue was carried out using preparative HPLC. Appropriate fractions were combined and concentrated in vacuo to give the title compound as an off-white solid (0.02 g, 55%): ¹H NMR (DMSO-d₆, 500 MHz) δ 10.54 (s, 1 H), 8.80 (t, 1 H, *J* = 5.5 Hz), 7.93 (d, 1 H, *J* = 7.9 Hz), 7.84 (s, 1 H), 7.64 (t, 1 H, *J* = 7.9 Hz), 7.50 (d, 1 H, *J* = 7.9 Hz), 6.18 (s, 1 H), 3.16 (m, 2 H), 2.09 (s, 3 H), 1.03 (t, 3 H, *J* = 7.1 Hz); LC/MS (ES+) 367 (M+H⁺); 399 (M+Na⁺).

Example 143**5-Cyano-3,6-dihydro-4-methyl-6-(3-(5-oxa-4-trifluoromethyl-2,3-diazole)phenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide**

Following the procedure of Method 10, the title compound was obtained from 6-(3-carboxyphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide, hydrazine and trifluoroacetic acid as a colorless solid in 42% yield: ¹H NMR (DMSO-d₆, 500 MHz) δ 10.54 (s, 1 H), 8.79 (t, 1 H, *J* = 5.5 Hz), 7.84 (d, 1 H, *J* = 7.7 Hz), 7.75 (s, 1 H), 7.57 (t, 1 H, *J* = 7.7 Hz), 7.51 (d, 1 H, *J* = 7.7 Hz), 6.13 (s, 1 H), 3.16 (m, 2 H), 2.09 (s, 3 H), 1.03 (t, 3 H, *J* = 7.1 Hz); ¹⁹F NMR (DMSO-d₆, 500 MHz) δ -73.35.

Example 144**6-(2-Bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine**

Following the procedure of Method 1, 6-(2-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine was obtained from 2-bromobenzaldehyde, acetoactamide and urea as a light yellow solid in 85% yield: ¹H NMR (DMSO-d₆,

400 MHz) δ 9.64 (s, 1 H), 7.81 (s, 1 H), 7.63 (d, 1 H, $J = 7.9$ Hz), 7.44 (m, 2 H), 7.29 (m, 1 H), 5.51 (s, 1 H), 2.01 (s, 3 H).

6-(2-(4-Chlorophenyl)phenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine

Following the procedure of Method 5, 6-(2-(4-chlorophenyl)phenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine was obtained from 6-(2-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine and 4-chlorophenylboronic acid as a colorless solid in 53% yield.

6-(2-(4-Chlorophenyl)phenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide

Following the procedure of Method 4, the title compound was obtained from 6-(2-(4-chlorophenyl)phenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine and ethyl isocyanate as a colorless solid in 47% yield: ^1H NMR (MeOH- d_4 , 400 MHz) δ 7.47 (m, 2 H), 7.35 (m, 5 H), 7.12 (d, 1 H, $J = 6.6$ Hz), 6.17 (s, 1 H), 3.16 (m, 2 H), 1.95 (s, 3 H), 1.05 (m, 3 H, $J = 7.0$ Hz); LC/MS (ES+) 395 ($M+H^+$).

Example 145

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(2-(3-pyridyl)phenyl)-(2H)-pyrimidine

Following the procedure of Method 5, 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(2-(3-pyridyl)phenyl)-(2H)-pyrimidine was obtained from 6-(2-bromophenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-(2H)-pyrimidine and pyridine-3-boronic acid, 1,3-propanediol cyclic ester as a colorless viscous oil in 29% yield: ^1H NMR (MeOH- d_4 , 400 MHz) δ 8.44 (m, 2 H), 7.74 (m, 1 H), 7.45 (m, 4 H), 7.14 (d, 1 H, $J = 7.5$ Hz), 5.12 (s, 1 H), 1.88 (d, 3 H, $J = 1.3$ Hz).

5-Cyano-3,6-dihydro-4-methyl-2-oxo-6-(2-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide


Following the procedure of Method 4, the TFA salt of the title compound was obtained from 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(2-(3-pyridyl)phenyl)-(2H)-

pyrimidine and ethyl isocyanate as a light yellow solid in 53% yield: LC/MS (ES+)
362 (M+H⁺).

Claims

What is claimed is:

1. A compound of formula IA



1A

its enantiomers, diastereomers, pharmaceutically acceptable salts, prodrugs and solvates thereof, wherein

10 R_1 is selected from the group consisting of hydrogen, alkyl and cycloalkyl;

Y_1 , Y_2 , Y_3 , Y_4 and Y_5 are independently selected from the group consisting of H, halogen, lower alkyl, O alkyl, O aryl, NH alkyl, NH aryl, N alkyl alkyl, N alkyl aryl, N aryl aryl with the proviso that at least one of Y_1 , Y_2 , Y_3 , Y_4 and Y_5 is aryl or heteroaryl;

15 R_4 is selected from the group consisting of alkyl, arylalkyl, cycloalkylalkyl, heteroarylalkyl, heterocycloalkylalkyl, CN, COR_5 , CO_2R_5 and $CONR_5R_6$;

R_5 and R_6 are independently selected from the group consisting of H, alkyl, haloalkyl, cycloalkyl, aminoalkyl, arylalkyl, cycloalkylalkyl, heteroarylalkyl, and heterocycloalkylalkyl;

20 Z is selected from the group consisting of O, S and NR_8 ;

R_8 is selected from the group consisting of H, CN, sulfonamido, OR_7 , alkyl, cycloalkyl, aryl, arylalkyl, heterocyclyl, heteroaryl and heteroarylalkyl; and

R_7 is selected from the group consisting of H, alkyl, arylalkyl, cycloalkylalkyl, heterocycloalkylalkyl, and heteroarylalkyl.

25

2. The compound of claim 1 wherein Y_2 is aryl or heteroaryl.

3. The compound of claim 2 wherein Y_2 is an optionally substituted phenyl, pyridyl, thiazolyl, pyrrolyl, or diazolyl.
4. The compound of claim 3 wherein said substituent is halo, alkyl, or haloalkyl.
5. The compound of claim 4 wherein said substituent is Br, Cl, F, methyl, ethyl, or CF_3 .
6. The compound of claim 1 wherein R_4 is alkyl, arylalkyl, CO_2R_5 or $CONR_5R_6$.
7. The compound of claim 1 wherein R_1 is alkyl; R_4 is selected from the group consisting of alkyl, arylalkyl, CO_2R_5 , and $CONR_5R_6$; R_5 and R_6 are independently selected from the group consisting of H, alkyl, haloalkyl, cycloalkyl, and arylalkyl; Z is selected from the group consisting of O, S, and NR_8 ; R_8 is selected from the group consisting of H and CN.
8. The compound of claim 1 wherein R_4 is CO_2R_5 and Z is O.
9. The compound of claim 1 wherein R_4 is $CONR_5R_6$ and Z is O.
10. The compound of claim 1 wherein R_4 is selected from the group consisting of alkyl and arylalkyl, and Z is O.
11. The compound of claim 1 wherein R_1 is CH_3 ; R_4 is CO_2R_5 ; R_5 is alkyl; and Z is O.
12. A compound selected from the group consisting of:
6-(3-biphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester;
5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-thienyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl ester;

- 6-(3-biphenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 6-(3-(4-chlorophenyl)phenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 5 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-thienyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-thienyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 10 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-(2,2,2-trifluoro)ethyl amide;
- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-isopropyl amide;
- 15 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(4-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide;
- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-(2,2,2-trifluoro)ethyl amide;
- 20 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-isopropyl amide;
- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide;
- 25 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclobutyl amide;
- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopentyl amide;
- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropylmethyl amide;
- 30 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;

- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(3-(2-pyrrole)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 5-cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 5 5-cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-(2,2,2-trifluoro)ethyl amide;
- 5-cyano-3,6-dihydro-6-(4-fluoro-3-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide;
- 5-cyano-3,6-dihydro-6-(2-fluoro-5-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 10 5-cyano-3,6-dihydro-6-(2-fluoro-5-(3-pyridyl)phenyl)-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide;
- 5-cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-3-pyridyl)phenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 15 5-cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-3-pyridyl)phenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-cyclopropyl amide;
- 5-cyano-3,6-dihydro-4-methyl-6-(3-(4-methyl-5-oxa-2,3-diazole)phenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 5-cyano-3,6-dihydro-4-methyl-6-(3-(5-oxa-4-trifluoromethyl-2,3-diazole)phenyl)-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide;
- 20 6-(2-(4-chlorophenyl)phenyl)-5-cyano-3,6-dihydro-4-methyl-2-oxo-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide; and
- 5-cyano-3,6-dihydro-4-methyl-2-oxo-6-(2-(3-pyridyl)phenyl)-1-(2H)-pyrimidinecarboxylic acid, 1-ethyl amide.
- 25
13. A method for treating a proliferative disease via inducing mitotic arrest comprising administering to a mammalian species an effective amount of a compound of claim 1.
- 30 14. The method according to claim 13 further comprising administering to the patient at least one other anti-cancer agent in combination with said compound.

15. The method according to claim 14 wherein said anticancer agent is cisplatin, doxorubicin, topoisomerase II inhibitors, etoposide, topoisomerase I inhibitors, CPT-11, topotecan, paclitaxel, docetaxel, epothilone, tamoxifen, thymidilate synthase inhibitors, and anti-metabolites.

5

16. The method according to claim 11 wherein the proliferative disease is cancer.

17. A method for treating a disease associated with the Eg5 motor protein
10 comprising administering to a mammalian species in need of such treatment an effective amount of a compound of claim 1.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/09494

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 239/42, 409/04, 401/04, 403/12; A61K 31/505, 31/506; A61P 9/14.
US CL : 514/274; 544/315, 318.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
U.S. : 514/274; 544/315, 318.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS ONLINE, EAST (IS&R)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00/78731 A1 (CELL TECH CHIROSCIENCE LIMITED) 28 December 2000 (28.12.00), especially page 60, Examples 31 & 32.	1-17
A	KAPPE, C. O. et. al. Synthesis and Reactions of "Biginelli-Compounds". Part I. Journal of Heterocyclic Chemistry, January-February 1989, Vol. 26, No. 1, pages 55-64, especially compound #27 on page 59.	1-17
A	REHWALD, M. et. al. Synthesis of Thieno[2,3-d]pyrimidines and Amino-pyrimidines from 2-Alkoxy-5-cyano-thioxopyrimidine Intermediates. Heterocycles. June 1998, Vol. 48, No. 6, pages 1157-1167, especially compound #6 on page 1159.	1-17
A	AGARWAL, N. et. al. Suitably Functionalised Pyrimidines as Potential Antimycotic Agents. Bioorganic & Medicinal Chemistry Letters. 2000, Vol. 10, pages 703-706, especially page 705.	1-17
A	Chem. Abstr., Vol. 114, No. 9. RAM, V. J. et. al. 'Chemotherapeutic Agents. XI. Synthesis of Pyrimidines and Azolopyrimidines as Leishmanicides'. 04 March 1991, Abstract No. 114: 81744t.	1-17

☒ Further documents are listed in the continuation of Box C.

See patent family annex.

Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

22 June 2002 (22.06.2002)

Date of mailing of the international search report

31 JUL 2002

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Mukund Shah

Telephone No. 703-308-1235

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/09494

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Chem. Abstr., Vol. 127, No. 12. KLEIDERNIGG, O. P. et. al. 'Synthesis and Reactions of Biginelli Compounds. 8. Separation of enantiomers of 4-Aryldihydropyrimidines by Direct Enantioselective HPLC. A Critical Comparison of Chiral Stationary Phases'. 22 September 1997, page 664, Abstract No. 127: 161790p.	1-17
A	Chem. Abstr., Vol. 133, No. 7. AL-AFALEQ et. al. 'Heterocyclic Synthesis via Enaminonitriles: An Efficient, One Step Synthesis of Some Novel Azolo[1,5-a]pyrimidine, Pyrimido[1,2-a]benzimidazole, Pyrido[1,2-a]benzimidazole, Pyrimidine and Pyrazole Derivatives'. 14 August 2000, pages 671-672, Abstract No. 133: 89496b.	1-17
A	Chem. Abstr., Vol. 129, No. 15. REHWALD, M. et. al. 'Synthesis of Thieno[2,3-d]pyrimidines and Aminopyrimidines from 2-Alkoxy-5-cyano-4-thioxopyrimidine Intermediates'. 12 October 1998, page 662, Abstract No. 129: 189296x.	1-17

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☒ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.